

VITAMIN B₂ (RIBOFLAVIN) CONTENT IN CEREAL PRODUCTS

S. Škrovánková, P. Sikorová

Received: July 8, 2010

Abstract

ŠKROVÁNKOVÁ, S., SIKOROVÁ, P.: *Vitamin B₂ (riboflavin) content in cereal products*. Acta univ. agric. et silvic. Mendel. Brun., 2010, LVIII, No. 5, pp. 377–382

Vitamin B₂ (riboflavin) is a water-soluble essential vitamin. Nowadays an increased risk for riboflavin deficiency may be seen in people on special diets (*diabetes mellitus*), smokers or heavy alcohol drinkers. In the Czech diet the main sources of the vitamin intake are milk and dairy products followed by cereals and meat. Cereals are good source of this vitamin as it is widely and regularly consumed in different forms. Analyses of the vitamin B₂ content in different types of cereal products (flours, breads, pastries, breakfast cereals, cooked pasta) of Czech origin using HPLC with reversed phase and UV detection were done. The vitamin B₂ content of chosen cereal products decreased in this progression: enriched wholemeal breakfast cereals (the best source of the vitamin), enriched wheat flours, breads – rye and wholemeal wheat breads, whole wheat and spelt flours, wheat bread, cooked whole wheat and rye spaghetti, wheat and multigrain pastries and finally scoured wheat flours.

vitamin B₂, riboflavin, cereal products, HPLC/UV

Vitamin B₂ – riboflavin is a water-soluble essential vitamin which is needed in the metabolism of amino acids, fatty acids, and saccharides. It also activates vitamin B₆ (pyridoxine) and folic acid and helps to create niacin. Riboflavin exerts its biological function through two coenzymes – flavine adenine dinucleotide (FAD) and flavine mononucleotide (FMN) which participate in oxidation-reduction reactions in numerous metabolic pathways exemplified by the oxidation of reduced NADH, the oxidation in the tricarboxylic acid cycle, and in the β-oxidation of fatty acids (Ball, 2006).

Humans cannot synthesize this vitamin and thus it must be obtained from their diet through absorption in the small intestine. The recommended daily intake (RDI) of riboflavin, related to protein and energetic intake, is from 1.2 to 1.7 mg per day in the Czech Republic (Velíšek, 2009).

Nowadays an increased risk for riboflavin deficiency may be seen in people on special diets (*diabetes mellitus*, peptic ulcer disease, weight loss), smokers, heavy alcohol drinkers and women using certain types of birth control. Therefore the solution could be the diet enriched with some of common vitamin sources as products with whole cereals

or riboflavin enriched products, e.g. bread and pastries. These products could be enriched by riboflavin (E 101) up to 50% of RDI, cereal mixtures and pasta even up to 100% of RDI. Vitamin fortification of foods is referred in Regulation No. 446/2004. This vitamin can be also used as a colorant which is referred in Regulation No. 304/2004.

Great natural food sources of vitamin B₂ include: entrails – liver, nuts, yeast, cheese, eggs, milk, lean meat and wholemeal cereals. In the Czech diet the main sources of the vitamin intake are milk and dairy products followed by cereals and meat (Hlúbík and Opltová, 2004). Cereals are good source of this vitamin as it is widely and regularly consumed in different forms – flour for home culinary purposes and technological production of cereal products, bread, pastries, breakfast cereals, pastas, and biscuits, especially for common cereals – wheat and rye.

The aim of this work was therefore to determine riboflavin content in common cereal products of the Czech origin.

Riboflavin is stable, not destroyed by heat or oxidation, unless exposed to ultraviolet radiation (420–560 nm). So it is a highly photosensitive compound. Knowledge of its photochemical behaviour

in aqueous solution over an appropriate pH range is needed to predict the shelf-life. The choice of the optimum pH is crucial for liquid preparations. Ahmad *et al.* (2004) determined that riboflavin solutions are most stable to UV and visible radiations at pH 5–6. Riboflavin content could be also decreased according to character of foods, technological and culinary processes (Lešková *et al.*, 2006), e.g. scouring and dissolution in cooking water (Al-Khalifa and Dawood, 1993).

Vitamins (vitamin B₂) in cereals occur especially in germ and aleuronic layer, so losses are connected to cereal scouring, primarily its level (Hägg and Kum-pulainen, 1994). Scoured cereal products without bran then contain up to ten times less B-complex vitamins than bran products. Losses of riboflavin due to baking are quite small (up to 10%). The significant losses of this vitamin (35–55%) are in cooked products due to leaching into water (Ayranci and Kaya, 1993; Velíšek, 2009). Freezing doesn't significantly affect riboflavin contents in breads (Hägg and Kum-pulainen, 1994).

In foodstuffs, the vitamin B₂ may be presented in free and phosphorylated forms, tightly bound but non-covalently to proteins. For vitamin extraction there can be used an acid hydrolysis or an enzymatic treatment (*takadiastasa*, *trypsin*, *clarasa*) (Ndaw *et al.*, 2000).

For vitamin determination several methods have been reported – microbiological assay (Golbach *et al.*, 2007), electrochemical method and High Performance Liquid Chromatography (Arella *et al.*, 1996, Tang *et al.*, 2006). The most widely used method in the determination of vitamin B₂ is HPLC using reverse phase with fluorescence (Andrés-Lacueva *et al.*, 1998; Cataldi *et al.*, 2003) and UV (Höller *et al.*, 2003) detection.

MATERIAL AND METHODS

Samples

20 cereal products are from the Czech production purchased in food markets.

- flours – wheat (bread, standard, pastry) flours
 - whole wheat and spelt flours
 - enriched wheat (bread, standard, pastry) flours – declared amount of riboflavin content of all enriched flours: 0.640 mg.100 g⁻¹
- breads: wheat, wholemeal (2 types), rye bread
- breakfast cereals: enriched wholemeal (3 types)
- pastries: wheat, multigrain (2 types) pastries
- cooked spaghetti: whole wheat and rye spaghetti (cooked in water at 90 °C for 10 min).

Reagents

- Calibration: riboflavin was obtained from Supelco (USA)
 - standard solutions: 0.25–4 µg.ml⁻¹

- extraction: HCl, trichloroacetic acid, Carrez I and Carrez II were obtained from P. Lukeš (Czech Republic)
- HPLC determination: CH₃COONa was purchased from P. Lukeš (Czech Republic), methanol of HPLC grade from Sigma – Aldrich Chemicals (Germany).

Vitamin extraction

- 100 ml 0.2 M HCl and 3 ml CCl₃COOH
- 30 ml of Carrez I and Carrez II.

Chromatographic conditions

- HPLC (Hewlett Packard 1100, USA)
- injection volume 20 µl
- mobile phase: gradient technique.

CH₃COONa : CH₃OH 87:13

- column SUPELCOSIL – LC8 (15 cm × 4.6 mm; 5 µm, Supelco, USA)
- column temperature 30 °C
- flow rate: 0.8 ml.min⁻¹
- Detector: UV/VIS DAD G1315A

λ = 270 nm.

Riboflavin Extraction and HPLC Determination

By reason of complicated matrices based on protein-polysaccharides, the first step was isolation of vitamin B₂ by acid hydrolysis. That is essential to denature the proteins and to release the vitamin from his association with the proteins and when the foodstuffs contain a lot of starch, to convert it into soluble form (Ndaw *et al.*, 2000).

Extraction was done with 0.2 M hydrochloric acid in a water bath at 95 °C for 1 hour with addition of 80% trichloroacetic acid in 50 minute. Moreover the proteins were subsequently precipitated by solutions Carrez I and Carrez II followed by filtration. All samples were kept away from light because of its low photostability.

The suitable conditions for the determination by HPLC technique were established partly according to Buňka *et al.* (2003). Mobile phase composed of sodium acetate, pH adjusted to 4.5 as riboflavin solutions are most stable to UV and visible radiations at pH about 5–6, and methanol, in the ratio 87:13 (v/v) with gradient technique (till 15 min., then 0:100) for 30 min. and flow rate 0.8 ml.min⁻¹.

For riboflavin detection UV/VIS DAD detector was used at the wavelength of 270 nm with ChemStation-Instrument 1 programme for the content calculation.

Method validation

Confirmatory analysis of validation parameters were experimentally verified and confirmed, as previously published by Buňka *et al.* (2003).

Calibration curve ($y = ax + b$): $y = 79.572x - 2.243$

Correlation coefficient: $r = 0.9995$

Recovery: $90.1 \pm 2.3\%$ – $91.4 \pm 3.1\%$ (known amount of vitamin B₂ was added to cereal samples)

Inter and intra-day repeatability: $SD < 3\%$.

RESULTS AND DISCUSSION

Table I. presents the content of riboflavin in 20 cereal products – flours, breads, pastries, breakfast cereals and cooked spaghetti. There is also shown contribution of 100 g cereal product to RDI (Recommended Daily Intake in the Czech Republic is 1.5 mg per day).

The vitamin B₂ content of chosen Czech cereal products decreased in this progression: enriched wholemeal breakfast cereals, enriched wheat flours, breads – rye and wholemeal wheat breads, whole wheat and spelt flours, wheat bread, cooked whole wheat and rye spaghetti, wheat and multigrain pastries and finally scoured wheat flours.

From the chosen products the best ones as for riboflavin content were wholemeal breakfast cereals ($0.553\text{--}2.779\text{ mg}\cdot 100\text{ g}^{-1}$) enriched with vitamins. This range represents 36.9–185.3% contribution to RDI for riboflavin. The results are in agreement with the literature (Hägg and Kumpulainen, 1993). Bran cereals also contain more vitamin B₂ than scoured ones as it was previously referred (Hägg and Kumpulainen, 1994). These products don't need any further culinary treatment so they are really good source of this vitamin.

Also all enriched wheat flours – bread (Fig. 1A), pastry and standard ($0.525\text{--}0.800\text{ mg}\cdot 100\text{ g}^{-1}$; contribution of 35.0–53.3% to RDI) and whole spelt and wheat flours (0.204 and $0.265\text{ mg}\cdot 100\text{ g}^{-1}$) seem as good sources. The vitamin concentration in scoured flours was very low (up to $0.087\text{ mg}\cdot 100\text{ g}^{-1}$ in wheat bread flour (Fig. 1B)), as expected. The results correspond to the producer values and are also in agreement with the literature (Batifoulouier *et al.*, 2005; Hägg and Kumpulainen, 1994; Ndaw *et al.*, 2000). The variability of the content in flours could be explained by milling procedure, genetic variability and growing location. So milling and the type of the flour (bread, standard, pastry) are very important for the responsibility in the differences of the vitamin results. And flours before consumption need another technological or culinary treatment so it could lead to another loss of vitamin due to that preparation.


Riboflavin content in bread products was higher than in flours. The concentration increased during bread making, suggesting a potential synthesis or contribution by yeast as referred Batifoulouier *et al.* (2005) and Batifoulouier *et al.* (2006).

Lower vitamin concentration was found in common wheat bread ($0.208\text{ mg}\cdot 100\text{ g}^{-1}$) than in wholemeal and rye breads ($0.322\text{--}0.561\text{ mg}\cdot 100\text{ g}^{-1}$; contribution of 21.5–37.4% to RDI). This could be related to the flour scouring level as the vitamin is contained especially in the bran. It reflects a more dras-

I: Riboflavin content in cereal products (meanSE) [$\text{mg}\cdot 100\text{ g}^{-1}$] and contribution of 100 g product to RDI [%]

Cereal products	Average riboflavin content	Contribution of 100 g product to RDI
Flour: Enriched wheat bread	0.800 ± 0.004	53.3
Enriched wheat pastry	0.584 ± 0.006	38.9
Enriched wheat standard	0.525 ± 0.005	35.0
Whole wheat	0.265 ± 0.005	17.7
Whole spelt	0.204 ± 0.004	13.6
Wheat bread	0.087 ± 0.006	5.8
Wheat pastry and standard	nd.	-
Bread: Rye	0.561 ± 0.010	37.4
Wholemeal 1	0.504 ± 0.013	33.6
Wholemeal 2	0.322 ± 0.013	21.5
Wheat	0.208 ± 0.008	13.9
Pastries: Wheat	0.094 ± 0.007	6.3
Multigrain 1	0.081 ± 0.007	5.4
Multigrain 2	nd.	-
Wholemeal breakfast cereals: 1	2.779 ± 0.007	185.3
2	1.246 ± 0.010	83.1
3	0.553 ± 0.012	36.9
Cooked spaghetti: Whole wheat	0.179 ± 0.007	11.9
Whole rye	0.223 ± 0.010	14.9

nd. – not detected



1: Chromatographic analysis of vitamin B₂ in enriched wheat bread flour (A) and wheat bread flour (B) by HPLC/UV

tic milling effect with refined flour than with unrefined, whole flour (Hägg and Kumpulainen, 1994).

Low concentrations of riboflavin in cooked spaghetti (0.179–0.223 mg.100 g⁻¹) of Czech origin, though wholemeal product, are due to vitamin leaching into water which correspond to results of Ayranci and Kaya (1993). According to them the percentage of riboflavin loss for macaroni during cooking at 90 °C was 18.3%, (10 min cooking with water) and 53.3% (20 min cooking). As expected, the lowest vitamin B₂ levels were found in pastries (up

to 0.094 mg.100 g⁻¹) made from scoured flours. So, common wheat or multigrain pastries are not very good source of this vitamin in the Czech Republic.

Riboflavin is water-soluble vitamin which leaches into water, e.g. during cooking with water. Other losses could happen during storage e.g. UV radiation impact. In some papers there have been proved that riboflavin losses depend on technological condition (Batifoulier *et al.*, 2005; Batifoulier *et al.*, 2006; Hägg and Kumpulainen, 1994; Prodanov *et al.*, 2004).

SUMMARY

Riboflavin is a water-soluble essential vitamin which is needed in the metabolism of amino acids, fatty acids, and saccharides (RDI=1.5 mg per day). Nowadays an increased risk for riboflavin deficiency may be seen in people on special diets (e.g. *diabetes mellitus*). In the Czech diet the main sources of the vitamin intake are milk and dairy products followed by cereals and meat. Cereals are good source of this vitamin as it is widely and regularly consumed in different forms. The aim of this work was therefore to determine riboflavin content in common cereal products of the Czech origin.

One of widely used method in the determination of vitamin B₂ is HPLC using reverse phase with UV detection. The vitamin B₂ content of chosen Czech cereal products, determined by this method, decreased in this progression: enriched wholemeal breakfast cereals (0.553–2.779 mg.100 g⁻¹; contribution of 36.9–185.3 % to RDI), which are really good source of the vitamin as they don't need any further culinary treatment; then enriched flours (0.525–0.800 mg.100 g⁻¹), breads – wholemeal with higher content as vitamin (0.322–0.561 mg.100 g⁻¹) which occurs especially in grain germ and aleuronic layer; following by wheat bread (0.208 mg.100 g⁻¹). In breads, in comparison to flours, could the riboflavin content increased during bread making (synthesis or by yeast contribution); next wheat flours – due to the cereal scouring the concentration was very low (up to 0.087 mg.100 g⁻¹; contribution of 5.8 % to RDI). Low concentrations of riboflavin in cooked spaghetti (0.179–0.223 mg.100 g⁻¹, contribution of 11.9–14.9 % to RDI) are due to vitamin leaching into water. The lowest vitamin B₂ levels were found in pastries (up to 0.094 mg.100 g⁻¹), so common wheat or multigrain pastries from scoured flours are not very good source of this vitamin.

SOUHRN

Obsah vitamínu B₂ (riboflavinu) v cereálních produktech

Riboflavin je ve vodě rozpustný vitamin, který je zapotřebí pro metabolismus aminokyselin, tuků a sacharidů (DDD = 1,5 mg na den). V dnešní době mají někteří lidé (např. nemocní s *diabetes mellitus*) větší potřebu přísunu vitamínu B₂. Zdroje tohoto vitamínu u lidí v České republice pochází především z mléka a mléčných výrobků, cereálií a masa. K nejvýznamnějším rostlinným zdrojům těchto vitaminů patří obiloviny kvůli jejich běžné a poměrně rozsáhlé konzumaci v různé formě. Cílem této práce proto bylo stanovení obsahu riboflavinu v běžných cereálních výrobcích českého původu. Jednou z nejvíce rozšířených metod pro jeho stanovení je HPLC s reverzní fází a UV detekcí. Obsah vitamínu B₂ ve vybraných cereálních produktech, stanovený výše uvedenou metodou, klesal v tomto pořadí: obohacené celozrnné snídaňové cereálie (0,553–2,779 mg.100 g⁻¹; kde 100 g výrobku zabezpečí 36,9–185,3 % DDD riboflavinu), které jsou tedy nejlepším zdrojem z hodnocených produktů bez nutnosti dalších kulinárních úprav. Následují vitaminem obohacené mouky (0,525–0,800 mg.100 g⁻¹), dále chleba – celozrnný s vyšším obsahem vitamínu (0,322–0,561 mg.100 g⁻¹), který se nachází především v klíčku a aleuronové vrstvě, a pšeničný chleba (0,208 mg.100 g⁻¹). Chleba v porovnání s odpovídajícími moukami může mít vyšší obsah vitamínu B₂ z důvodu nárůstu v průběhu pečení (syntéza nebo přídavek droždí). Bílé pšeničné mouky měly poměrně nízký obsah riboflavinu s ohledem na vymílání těchto mouk (do 0,087 mg.100 g⁻¹; 5,8 % DDD). Nízké obsahy riboflavinu ve vařených špagetách (0,179–0,223 mg.100 g⁻¹, 11,9–14,9 % DDD) souvisí s vyluhováním do vodného prostředí. Nejnížší hodnoty obsahů vitamínu B₂ byly zjištěny v pečivu (do 0,094 mg.100 g⁻¹), tedy běžné vymílané bílé a vícezrnné pečivo není výhodným zdrojem tohoto vitamínu.

vitamin B₂, riboflavin, cereální produkty, HPLC/UV

REFERENCES

- AHMAD, I., FASIHULLAH, Q., NOOR, A., ANSARI, I. A., MANZAR ALI, Q. N., 2004: Photolysis of riboflavin in aqueous solution: a kinetic study. *International Journal of Pharmaceutics*, 280, 1–2: 199–208. ISSN 0378-5173.
- AL-KHALIFA, A. S., DAWOOD, A. A., 1993: Effects of cooking methods on thiamine and riboflavin contents of chicken meat, *Food Chemistry*, 48, 1: 69–74. ISSN 0308-8146.
- ANDRÉS-LACUEVA, C., MATTIVI, F., TONON, D., 1998: Determination of riboflavin, flavin mononucleotide and flavin-adenine dinucleotide in wine and other beverages by high-performance liquid chromatography with fluorescence detection. *Journal of Chromatography A*, 823, 1–2: 355–363. ISSN 0021-9673.
- ARELLA, F., LAHÉLY, S., BOURGUIGNON, J. B., HASSELMANN, C., 1996: Liquid chromatographic determination of vitamins B₁ and B₂ in foods: a collaborative study. *Food Chemistry*, 56, 1: 81–86. ISSN 0308-8146.
- AYRANCI, G. AND S. KAYA, S., 1993: Kinetic analysis of the loss of some B-vitamins during the cooking of macaroni. *Nahrung*, 37, 2: 153–155. ISSN 0027-769X.
- BALL, G. F. M., 2006: *Vitamins In Foods: Analysis, Bioavailability, and Stability*. 1. vyd. Boca Raton: CRC Press, 785 s. ISBN 978-1-57444-804-7.
- BATIFOULIER, F., VERNY, M.-A., CHANLIAUD, E., RÉMÉSY, C., DEMIGNÉ, C., 2005: Effect of different breadmaking methods on thiamine, riboflavin and pyridoxine contents of wheat bread. *Journal of Cereal Science*, 42, 1: 101–108. ISSN 0733-5210.
- BATIFOULIER, F., VERNY, M.-A., CHANLIAUD, E., RÉMÉSY, C., DEMIGNÉ, C., 2006: Variability of B vitamin concentrations in wheat grain, milling fractions and bread products. *European Journal of Agronomy*, 25, 2: 163–169. ISSN 1161-0301.
- BUŇKA, F., SEVEROVÁ, M., HRABĚ, J., 2003: Vliv sterilace na změny riboflavinu v tavených sýrech

- určených do bojových dávek potravin. In: Sborník VVŠ PV, 2, 121–130.
- CATALDI, T. R. I., NARDIELLO, D., CARRARA, V., CIRIELLO, R., DE BENEDETTO, G. E., 2003: Assessment of riboflavin and flavin content in common food samples by capillary electrophoresis with laser-induced fluorescence detection. *Food Chemistry*, 82, 2: 309–314. ISSN 0308-8146.
- GOLBACH, J. L., CHALOVA, V. I., WOODWARD, C. L., RICKE, S. C., 2007: Adaptation of *Lactobacillus rhamnosus* riboflavin assay to microtiter plates. *Journal of Food Composition and Analysis*, 20, 7: 568–574. ISSN 0889-1575.
- HÄGG, M., KUMPULAINEN, J., 1993: Thiamine and Riboflavin Contents in Domestic and Imported Cereal Products in Finland. *Journal of Food Composition and Analysis*, 6, 4: 299–306. ISSN 0889-1575.
- HÄGG, M., KUMPULAINEN, J., 1994: Thiamine and Riboflavin Contents of Finnish Breads and Their Corresponding Flours. *Journal of Food Composition and Analysis*, 7, 1–2: 94–101. ISSN 0889-1575.
- HLÚBIK, P., OPLTOVÁ, L., 2004: *Vitaminy*. 1. vyd. Praha: Grada Publishing, 232 s. ISBN 80-247-0373-4.
- HÖLLER, U., BRODHAG, C., KNÖBEL, A., HOFMANN, P., SPITZER, V., 2003: Automated determination of selected water-soluble vitamins in tablets using a bench-top robotic system coupled to reversed-phase (RP-18) HPLC with UV detection. *Journal of Pharmaceutical and Biomedical Analysis*, 31, 1: 151–158. ISSN 0731-7085.
- LEŠKOVÁ, E., KUBÍKOVÁ, J., KOVÁČIKOVÁ, E., KOŠICKÁ, M., PORUBSKÁ, J., HOLČÍKOVÁ, K., 2006: Vitamin losses: Retention during heat treatment and continual changes expressed by mathematical models. *Journal of Food Composition and Analysis*, 19, 4: 252–276. ISSN 0889-1575.
- NDAW, S., BERGAENTZLÉ, M., AOUDÉ-WERNER, D., HASSELMANN, C., 2000: Extraction procedures for the liquid chromatographic determination of thiamin, riboflavin and vitamin B₆ in foodstuffs. *Food Chemistry*, 71, 1: 129–138. ISSN 0308-8146.
- PRODANOV, M., SIERRA, I., VIDAL-VALVERDE, C., 2004: Influence of soaking and cooking on the thiamin, riboflavin and niacin contents of legumes. *Food Chemistry*, 84, 2: 271–277. ISSN 0308-8146.
- REGULATION No. 446/2004; Sbírka Zákonů ČR, 2004, částka 159, 16.7.2004.
- REGULATION No. 304/2004; Sbírka Zákonů ČR, 2004, částka 100, 6.5.2004.
- TANG, X., CRONIN, D.A., BRUNTON, N.P., 2006: A simplified approach to the determination of thiamine and riboflavin in meats using reverse phase HPLC. *Journal of Food Composition and Analysis*, 19, 8: 831–837. ISSN 0889-1575.
- VELÍŠEK, J., 2009: *Chemie potravin I*. 3. vyd. Tábor: Osis, 602 s. ISBN 978-80-86659-15-2.

Address

Ing. Soňa Škrovánková, Ph.D., Ing. Pavlína Sikorová, Ústav biochemie a analýzy potravin, Fakulta technologická, Univerzita T. Bati ve Zlíně, nám. T. G. Masaryka 275, 762 72 Zlín, Česká republika, e-mail: skrovankova@ft.utb.cz