

WIND EROSION ON HEAVY-TEXTURED SOILS: CALCULATION AND MAPPING

J. Kozlovsky Dufková, J. Podhrázská

Received: September 5, 2011

Abstract

KOZLOVSKY DUFKOVÁ, J., PODHRÁZSKÁ, J.: *Wind erosion on heavy-textured soils: calculation and mapping*. Acta univ. agric. et silvic. Mendel. Brun., 2011, LIX, No. 6, pp. 199–208

The equation that expresses the influence of factors affecting soil aggregates breakdown, and thus wind erosion, originated from the results of laboratory simulations of soil aggregates breakdown due to low temperatures treatment, field measurements of air temperature and soil moisture, and statistical evaluation of gained outcomes. All the analyses, whether field or laboratory, were realized on three different soils from three different localities of the Bílé Karpaty Mountains foothills – Ostrožská Nová Ves, Blatnice pod Svatým Antonínkem, and Suchá Loz. The statistically significant factors, influencing the soil aggregates breakdown, were determined using multiple regression analysis and stepwise regression. Soil moisture content at time of freezing was the most significant factor affecting soil aggregates breakdown, content of soil particles < 0.01 mm was the least significant one. Based on the results of laboratory and field research there was created a map of heavy-textured soils that are vulnerable to wind erosion.

soil aggregates breakdown, non-erodible fraction, multiple regression analysis, stepwise regression, GIS

Wind erosion is a serious problem in many parts of the world and extensive aeolian deposits from past geologic eras give evidence that it is not a recent phenomenon. Area of 19% of agricultural soils of the Czech Republic is potentially vulnerable to wind erosion (www.sowac-gis.cz). Potential vulnerability to wind erosion relates only to light-textured soils whereas heavy-textured soils are not vulnerable to wind erosion due to the formation of clods and surface crust that resist the blast of wind (Chandler *et al.*, 2005; Langston and McKenna Neuman, 2005; Grešová and Středanský, 2011). However, heavy-textured soils can become as erodible as light-textured ones during winter period. This phenomenon occurs in the foothills of the Bílé Karpaty Mountains, east of the Czech Republic (Podhrázská *et al.*, 2008). Freezing and thawing and freeze-drying of the soil are the main reasons that make resistant heavy-textured soils vulnerable to soil loss by wind. The mentioned overwinter processes are the main contributing factors causing soil aggregates breakdown and subsequent wind erosion also on the Canadian Prairies and northern

Great Plains (Bullock *et al.*, 2001). However, the processes proceed here under different climatic conditions than in the Czech Republic due to more highly altitude and thus lower air temperature in the winter period.

The paper deals with the assessment of the most statistically important factors that influence the soil aggregates breakdown in the foothills of the Bílé Karpaty Mountains, and with the quantification of the relationship between the factors using equation. The paper also deals with the map processing to express the localization and size of the area of heavy-textured soils vulnerable to wind erosion.

Data of soil and meteorological factors are used for the surface and space identification and mapping of areas vulnerable to wind erosion. Similar maps were created by Cablík and Jůva (1963), Prettl (1963) and Stehlík (1983) in the past for the territory of former Czechoslovakia, recently by Janeček *et al.* (2000) or SOWAC GIS (2011) for the territory of the Czech Republic. All the mentioned maps delimitate the vulnerability to wind erosion only on light-textured soils in warm and dry territories. Presented paper

shows new maps that were created according to the results of the laboratory and field research on heavy-textured soils vulnerable to wind erosion during specific winter conditions.

MATERIALS AND METHODS

The equation that expresses the influence of factors affecting soil aggregates breakdown, and thus wind erosion, originated from the results of laboratory simulations of soil aggregates breakdown due to low temperatures treatment, field measurements of air temperature and soil moisture, and statistical evaluation of gained outcomes.

All the analyses, whether field or laboratory, were realized on three different soils from three different localities of the Bílé Karpaty Mountains foothills – Ostrožská Nová Ves, Blatnice pod Svatým Antonínkem, and Suchá Loz. Some of the general characteristics of the soils used in this study are given in Table I.

Own methodology which could simulate the impact of climatic conditions on soil aggregates breakdown was developed. The new methodology was created on the basis of literature survey that deals with simulations of low temperatures impact on soil aggregates breakdown and on the basis of analysis of each individual overwintering method shortcomings.

Laboratory analyses were done with the soil samples according to the methodology that is fully described by Kozlovský Dufková (2010a) or Kozlovský Dufková (2010b).

Soil samples from soil surface were air-dried and passed through dry aggregate size distribution (DASD) analysis (Hagen *et al.*, 1987) to determine the amount of non-erodible fraction (NEF, aggregates > 2 mm in diameter) (Švehlík, 1990). Many studies (e.g. Hinman and Bisal, 1968; Staricka and Benoit, 1995; Bullock *et al.*, 1999) have shown that soil water content at the time of freezing is an important factor influencing soil disruption. For this reason, water was carefully added to NEF and four soil moisture levels were established – near saturation (Θ_s), field capacity (Θ_{PK}), wilting point (Θ_v), and no water added (dried soil, Θ_0). NEF was consequently exposed to the effect of low temperatures on the laboratory conditions. Effects of freeze/thaw (FT) and freeze/dry (FD) processes were observed. Temperature cutoffs of $-2\text{ }^{\circ}\text{C}$ for freezing, $+2\text{ }^{\circ}\text{C}$ for thawing, and $-2\text{ }^{\circ}\text{C}$ for freeze-drying were used to estimate the number of FT and FD cycles. With regard to the

fact that FT and FD processes occur several times during winter, the cycles were repeated several times. DASD analysis was made after every cycle and percentage content of NEF was determined. NEF was repeatedly exposed to low temperatures in FT and FD processes to investigate how many cycles are required for disruption of all NEF.

Field measurements of air temperature and soil moisture were realized. The air temperature in 2.0 m above the ground was measured by Hobo sensor (Onset Computer Corporation), the ground air temperature at the soil surface was measured by Dallas DS18B20 sensor (Maxim/Dallas Semiconductor Products, Maxim Integrated Products). VIRRIB sensor (Amet) was measured soil surface moisture at 0.5 m below the ground.

The task of the temperature measurements was to record the number of FT/FD cycles in the field conditions during the winter period from November 2009 to April 2010 and from November 2010 to April 2011.

Results from laboratory simulations and field measurements were statistically evaluated to check whether the suggested methodology of soil aggregates breakdown in the laboratory conditions simulates the field FT/FD processes exactly. Statistical analyses were made using Minitab 15.1. (Minitab Inc.), Unistat 5.1 (Unistat Ltd.) and MS Office Excel programmes.

For the graphical representation of the area of heavy-textured soils vulnerable to wind erosion there was used the system of Evaluated Soil – Ecological Units (ESEU). This system includes basic information about the soils in the Czech Republic and consists of five-number code. Each number (or couple of numbers) expresses certain characteristics of soil. The system of ESEU then respects all attributes and dissimilarities of certain agronomical area (soil, climatic, and morphological characteristics). The area and spatial distribution of various types of ESEU is expressed through the digital file of ESEU maps, i.e. the boundary-lines of ESEU area with their numerical codes. This system is used e.g. for various GIS analyzes of soil characteristics in the Czech Republic according to the value of appropriate codes or their combination.

RESULTS AND DISCUSSION

Comparison of the field and laboratory simulated FT/FD processes was realized on the basis of comparison of erodible fraction cumulative

I: Pedological characteristics of soils from three experimental areas

Experimental area	Soil type	ESEU code	Soil particles < 0.01 mm (%)	Non-erodible particles > 2 mm (%)	Organic matter (%)	CaCO ₃ (%)
Ostrožská Nová Ves	chernozem	07	52.8	54.3	2.3	0.7
Blatnice pod Sv. Antonínkem	chernozem	06	58.7	58.3	2.9	0.2
Suchá Loz	cambisols	20	62.7	61.6	3.7	0.2

increases (EF; aggregates < 2 mm in diameter) that came after particular FT/FD cycles (Fig. 1). The reason of using cumulative increases of EF was the unequal content of NEF in the analysed samples at the beginning of the field and simulated FT/FD process. The cumulative increases show the gradual increasing in the amount of disintegrated soil aggregates from the first to the last cycle of field and simulated FT/FD processes.

Graphical comparison of field and simulated FT/FD processes is based on the subjective assessment. Objective comparison was made by the statistical analyses in Unistat and Minitab programmes using of:


- Pearson correlation coefficient that expresses dependence between EF cumulative increases of field and simulated FT/FD processes; the Pearson correlation coefficient R measures the extent to which two continuous variables are related (Minařík, 2007), $R = 0.9130$ – 0.9935 for this study; the P -value tells if the correlation coefficient is significantly different from 0, $P = 0.001$ – 0.053 for this study,
- two-sample t -test and confidence interval for field vs. simulated FT/FD processes; these procedures

are used to make inferences about the difference between two population means (Friesl, 2004); example of results for Blatnice pod Sv. Antonínkem study area (winter period of 2009–2010) are given in Fig. 2 as an individual value plot and a box plot.


Individual value plot is useful to look at side-by-side plots of the data in order to compare the groups. Individual value plots illustrate the dispersion (each dot represents a value observed in a sample) and means (the blue symbols on each plot represents the mean of the sample). A box plot can tell a lot about the distribution of the data. The box represents the middle 50% of the differences, the line through the box represents the median difference, lines extending from the box represent the upper and lower 25% of the differences (excluding outliers) (Hendl, 2004).

Statistical analyses show that the field and simulated FT/FD processes of soil aggregates breakdown do not significantly differ from 95%. It means that the suggested methodology of FT/FD processes in the laboratory conditions really simulates the field FT/FD processes.


Statistically significant factors that influence the soil aggregates breakdown were determined using



1: Effect of number of FT/FD cycles on cumulative increases of erodible fraction



2: Individual value plots and box plots for erodible fraction cumulative increases of FT (a, b) and FD (c, d) simulated and field processes for the soil from the study area of Blatnice pod Sv. Antonínkem, winter period of 2009–2010



3: Dendrogram displaying the groups formed by clustering of variables and their similarity levels

multiple regression analysis (by Minitab program) and stepwise regression (by Unistat program). Regression analysis investigates and models the relationship between a response (dependent variable) and predictors (independent variables). The result of regression analysis is a regression

model/equation. Stepwise regression removes and adds variables to the regression model for the purpose of identifying a useful subset of the predictors (Hendl, 2004).

Stepwise regression analysis was made using final content of NEF after several FT/FD cycles in

the winter period as the dependent variable. The independent variables included initial content of NEF before FT/FD processes effect, number of cycles of FT/FD processes in the winter period, soil moisture content at time of freezing, content of soil particles < 0.01 mm, content of organic matter, and content of CaCO_3 . Multicollinearity was detected for the variables content of organic matter and CaCO_3 during creating of correlation matrix (these two variables were highly correlated with other variables in the matrix), thus these variables were excluded from the regression model by Minitab and Unistat programmes. The resulting relationship is given by equation (1). Correlation coefficient for the relationship is 0.8940 and P-value = 0.000.

$$\begin{aligned} \text{NEF}_{\text{end}} = & -14.09 + 0.49\text{NEF}_{\text{start}} - 0.43\text{V} - 1.02\text{FT} - \\ & -1.34\text{FD} + 0.62\text{M}, \end{aligned} \quad (1)$$

where NEF_{end} = final content of non-erodible fraction after several FT/FD cycles in the winter period (%), $\text{NEF}_{\text{start}}$ = initial content of non-erodible fraction before FT/FD processes effects (%), FT = number of freeze-thawing cycles in the winter period, FD = number of freeze-drying cycles in the winter period, V = soil moisture content at time of freezing (%), M = content of soil particles < 0.01 mm (%).

According to the cluster analysis and dendrogram (by Minitab program, Fig. 3), the factor that affects the soil aggregates breakdown at most in the winter period is a variable “soil moisture content at time of freezing”. Variable “content of soil particles < 0.01 mm” influences the soil aggregates breakdown at least.

The tested soil types were within delimitation and mapping of ESEU submitted into three codes of main soil units (Table I). Chosen soil units were selected from the state-wide database of ESEU. Their spatial and surface representation was realized through the map of potential vulnerability of heavy-textured soils to wind erosion (Fig. 4–6).

CONCLUSIONS

The relationship (1) is able to determine the amount of NEF after several FT/FD cycles in the winter period, when soil moisture content at time of freezing and content of soil particles < 0.01 mm are known. The equation is not able to specify actual amount of eroded soil, however is able to tell if the heavy-textured soil is vulnerable to loss by wind after winter period on the basis of prediction for the erosion threshold of 40% NEF. Bisal and Ferguson (1968) reported that soils with < 40% NEF were prone to wind erosion and they called this value as an erosion threshold.

SUMMARY

Wind erosion, a phenomenon affecting light-textured soils primarily, occurs also on heavy-textured soils in some regions of the Czech Republic. The foothills of the Bílé Karpaty Mountains belong to the areas, where the anomaly could be found. The process of wind erosion affects the soil first of all in winter and pre-spring period, when a breakdown of soil aggregates can be observed due to low air temperatures.


Heavy-textured soils from three study sites in the foothills of the Bílé Karpaty Mountains (Ostrožská Nová Ves, Blatnice pod Svatým Antonínkem, and Suchá Loz), were subjects of physical soil analyses. The objective of the laboratory analyses/simulations was to determine the influence of overwinter processes on soil aggregates breakdown and thus determine the vulnerability to soil loss by wind. Two overwinter processes on the soil were observed – a freeze and thawing process (FT), and a freeze-drying process (FD), where the various numbers of cycles were used. The analyses were realized with soils of four water contents (near saturation, field capacity, wilting point, and dried soil). Simultaneously, field measurements of air temperature and soil moisture were realized whereas the results of the measurements should determine the number of FT and FD cycles directly in the field conditions.

The statistically significant factors, influencing the soil aggregates breakdown, were determined using multiple regression analysis and stepwise regression. The most significant factor was soil moisture content at time of freezing, the least significant factor, from the point of view of soil aggregates breakdown, was content of soil particles < 0.01 mm. The equation that expresses the influence of factors affecting soil aggregates breakdown, and thus wind erosion, originated from the results of laboratory simulations of soil aggregates breakdown due to low temperatures treatment, field measurements of air temperature and soil moisture, and statistical evaluation of gained outcomes.


Creating of the map of potential vulnerability of heavy-textured soils to wind erosion completed information about spatial and surface location of areas prone to wind erosion in the territory of the Czech Republic.

Acknowledgement


Supported by the Ministry of Agriculture of the Czech Republic, project No. MZE QH82099.



4: Heavy-textured soils vulnerable to wind erosion in South Moravian region



5: Heavy-textured soils vulnerable to wind erosion in Zlín region



6: Heavy-textured soils vulnerable to wind erosion in the Czech Republic

REFERENCES

- BULLOCK, M. S., LARNEY, F. J., IZAURRALDE, R. C. and FENG, Y., 2001: Overwinter changes in wind erodibility of clay loam soils in southern Alberta. *Soil Science Society of America Journal*, Vol. 65, p. 423–430.
- BULLOCK, M. S., LARNEY, F. J., MCGINN, S. M. and IZAURRALDE, R. C., 1999: Freeze-drying processes and wind erodibility of a clay loam soil in southern Alberta. *Canadian Journal of Soil Science*, Vol. 79(1), p. 127–135.
- CABLÍK, J. and JŮVA, K., 1963: *Protierozní ochrana půdy*. 1. vyd. Praha: SZN Praha, 324 s.
- CHANDLER, D. G., SAXTON, K. E. and BUSACCA, A. J., 2005: Predicting wind erodibility of loessial soils in the Pacific Northwest by particle sizing. *Arid Land Research and Management*, Vol. 19(1), p. 13–27.
- FRIESL, M., 2004: *Pravděpodobnost a statistika hypertextově*. Verze 2004-09-15 HTML. URL: <<http://home.zcu.cz/~friesl/hpsb/tit.html>> [cit. 2011-07-01].
- GREŠOVÁ, L. and STREĎANSKÝ, J., 2011: *Veterná erózia v krajine – súčasné trendy, metódy a spôsoby výpočtov*. 1. vyd. Nitra: SPU v Nitre, 108 p. ISBN 978-80-552-0572-4.
- HAGEN, L. J., SKIDMORE, E. L. and FRYREAR, D. W., 1987: Using two sieves to characterize soil aggregate size distribution. *Transactions of the American Society of Agricultural Engineers*, Vol. 30(1), p. 162–165.
- HENDL, J., 2004: *Přehled statistických metod zpracování dat*. 3. vyd. Praha: Portál, 696 p. ISBN 978-80-7367-482-3.
- HINMAN, W. C. and BISAL, F., 1968: Alterations of soil structure upon freezing and thawing and subsequent draining. *Canadian Journal of Soil Science*, Vol. 48, p. 193–197.
- JANEČEK, M., TIPPL, M., PIVCOVÁ, J. and VETIŠKOVÁ, D., 2000: *Způsoby omezení degradace půd erozí a systémy protierozní ochrany. Mapy potenciální ohroženosti zemědělských půd České republiky vodní a větrnou erozí*. Výstup z řešení projektu NAZV EP7057. Praha: VÚMOP Praha - Zbraslav.
- KOZLOVSKY DUFKOVÁ, J., 2010a: Assessment of influence of overwinter processes on soil degradation by wind erosion. In: *16th international congress of Soil and Water Conservation* (proceedings [CD-Rom]). Santiago de Chile: International Soil Conservation Organization, 8.–12. 11. 2010, p. 384–389.
- KOZLOVSKY DUFKOVÁ, J., 2010b: Vliv nízkých teplot na rozpad půdních částic těžkých půd. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis*, Vol. 58(2), p. 161–168. ISSN 1211-8516.
- LANGSTON, G. and MCKENNA NEUMAN, C., 2005: An experimental study on the susceptibility of crusted surfaces to wind erosion: A comparison of the strength properties of biotic and salt crusts. *Geomorphology*, Vol. 72, p. 40–53.
- MINAŘÍK, B., 2007: *Statistika II*. 1. vyd. Brno: MZLU v Brně, 136 p. ISBN 978-80-7375-033-6.
- PODHRÁZSKÁ, J., NOVOTNÝ, I., ROŽNOVSKÝ, J., HRADIL, M., TOMAN, F., DUFKOVÁ, J., MACKŮ, J., KREJČÍ, J., POKLADNÍKOVÁ, H. and STŘEDA, T., 2008: *Optimalizace funkcí větrolamů v zemědělské krajině*. Metodika. Praha: VÚMOP Praha, 81 p. 978-80-904027-1-3.
- PRETL, J., 1963: Mapa ohrožení zemědělských půd ČSSR větrnou erozí. *Vodní hospodářství*, č. 5, 165 s.
- STARICKA, J. A. and BENOIT, G. R., 1995: Freeze-drying effects on wet and dry soil aggregate stability. *Soil Science Society of America Journal*, Vol. 59, p. 218–223.
- STEHLÍK, O., 1983: *Potenciální eroze půdy v ČSR*. Mapa. Geografický ústav ČSAV Brno, Geodetický a kartografický podnik v Praze.
- ŠVEHLÍK, R., 1990: Hranice erodovatelnosti půdy větrem. *Geografický časopis*, Vol. 42(3), p. 309–319.
- www.sowac-gis.cz, 2011: URL: <http://ms.sowac-gis.cz/mapserv/dhtml_eroze/index.php?project=dhtml_eroze&> [cit. 2011-06-27].

Address

Ing. Jana Kozlovsky Dufková, Ph.D., Ústav aplikované a krajinné ekologie, Mendelova univerzita v Brně, Zemědělská 1, 613 00 Brno, Česká republika, Ing. Jana Podhrázská, Ph.D., Výzkumný ústav meliorací a ochrany půdy, v. v. i., oddělení Pozemkové úpravy a využití krajiny, Lidická 25/27, 602 00 Brno, Česká republika, e-mail: janadufkova@email.cz, podhrazska.jana@vumop.cz

