

EFFECT OF BARN AIRSPACE TEMPERATURE ON COMPOSITION AND TECHNOLOGICAL PARAMETERS OF BULK MILK PRODUCED BY DAIRY COWS OF CZECH FLECKVIEH AND HOLSTEIN BREEDS

O. Polák, D. Falta, O. Hanuš, G. Chládek

Received: August 31, 2011

Abstract

POLÁK, O., FALTA, D., HANUŠ, O., CHLÁDEK, G.: *Effect of barn airspace temperature on composition and technological parameters of bulk milk produced by dairy cows of Czech Fleckvieh and Holstein breeds.* Acta univ. agric. et silvic. Mendel. Brun., 2011, LIX, No. 6, pp. 271–280

On two farms (A and B), samples of bulk milk produced by dairy cows of the Holstein (H) and Czech Fleckvieh (CF) breeds were collected every week on the same day within the time interval of 36 weeks. The aim of this sampling was to determine the effect of barn airspace temperature on milk composition and its technological parameters. The following average values of all bulk milk samples ($n = 72$) were recorded: barn airspace temperature (BAT) 12.08 °C; rennet coagulation time (RCT) 211 sec.; curd quality class (CQC) 1.53; titratable acidity (TA) 7.16 °SH; protein content (P) 3.46 %; fat content (F) 4.09 %; lactose content (L) 4.79 %; and solids non fat (SNF) 8.86 %. As compared with H, dairy cows of CF breed produced milk with statistically significantly higher ($P < 0.01$) contents of P (+0.27 %) and SNF (+0.26 %). The remaining differences were statistically insignificant ($P > 0.05$): BAT (–2 °C); RCT (–4 sec.); CQC (0 class); TA (+0.11 °SH); F (+0.05 %) and L (–0.03 %). As far as the effect of temperature was concerned, calculated correlation coefficients and plotted graphs indicated a marked effect of BAT on RCT; F and P. This effect was in all cases markedly negative: at lower BAT, F and P values were higher and RCT longer while at higher temperatures they were lower and shorter. These trends were similar in both breeds regardless to differences in average values of aforementioned parameters.

barn airspace temperature, milk, composition, technological parameters, Holstein, Czech Fleckvieh

Although the milk cattle shows a high adaptability to a wide scale of climatic conditions, its performance can be influenced by great temperature fluctuations occurring within the year. Nowadays, effects of the heat stress represents a tropical problem also in Eastern and Central Europe. Summer climate causes the heat stress of dairy cows and the heat stress results in am depression in milk production. The heat stress occurs in situations when the ambient temperature is higher than that of the animal's thermal neutral zone (Novák *et al.*, 2009). According to Vokřálková and Novák (2005), the thermoneutral zone of dairy cows ranges from –5 to +24 °C.

Other authors reminded that in high-yielding (i.e. > 6,500 kg) and, especially, older cows, the thermal stress developed at temperatures > 21 °C (Novák *et al.*, 2009; Vokřálková and Novák, 2005). The heat stress problem is getting worse as production levels continue to rise (Mittlöhner *et al.*, 2002; Beatty *et al.*, 2006). The summer depression in production of milk causes significant economical losses in the dairy industry. The basic condition of dairy farm management depends on the knowledge of and understanding to factors affecting milk production at most, i.e. not only nutrition and health status of dairy cows but also the parity and calving season,

technological systems, and, above all, microclimatic conditions (Maust *et al.*, 1972; Gader *et al.*, 2007). Livestock performance is affected by heat stress mainly due to the fact that animals having problems with high temperatures and heat try to control their thermoregulation and heat production by reduced feed intake (Davis *et al.*, 2003; Mader *et al.*, 2004).

Within a species, the variation in milk composition and yield is dependent on many factors. Some of them are of genetic nature while others concern stage of lactation, daily variation, parity, type of feeding, udder health, and season (Haenlein, 2003). Climatic conditions are known as seasonal changes which influence the milk composition. There is a negative correlation between the environmental temperature on the one hand and amounts of milk fat and protein on the other. When the temperature is increasing the solids non fat tends to decrease (Ozrenk and Inci, 2008). Ng-Kwai-Hang *et al.* (1984) and Lacroix *et al.* (1996) reported that the percentage of fat, protein and casein was influenced by the seasonal variations. Hanuš *et al.* (2008) observed influence of summer period on milk composition particularly on protein and solids non fat which decreased. Also Dolejš *et al.* (1996) mentioned a decrease in protein and solid non fat content in milk with increase of air temperature. It is clear that influence of dairy cow milk yield level on fat content (Hanus *et al.*, 2007) is more intensive in Czech Fleckvieh but less intensive in Holstein (Janů *et al.*, 2007) which is comparable to influence of environmental temperature variation on fat. Dolejš *et al.* (1996) found also the milk fat content depression with air temperature increase. Sevi *et al.* (2001) found high ambient temperatures to adversely affect the yield and cheese-making parameters of milk by the clotting time and the rate of clot formation and decreasing clot firmness. The photoperiod (i.e. light-to-dark ratio) can also induce marked changes in milk yield and composition (Casati *et al.*, 1998). In fact, a high light-to-dark ratio leads to a reduction in fat and protein contents of milk, probably as a consequence of a greater secretion of prolactin whose concentration in plasma is higher in the summer than in the winter (Tucker, 1989). Lactation period moved forward progressing and when the environmental heat degree increased, the fat content decreased (Sekerden, 1999; Yetismeyen, 2000). Jōdu *et al.* (2008) concluded that an increase of protein content of milk has resulted in reducing rennet coagulation time of milk.

Marked environmental effects on milk composition and technological quality of milk also cannot be negligible. These effects are usually involved into such models as effect of breed, year or season. For example De Marchi *et al.* (2007) mentioned that in their experiments, milk producing by dairy cows of Holstein-Friesian breed showed the worst coagulation (including RCT) among all other breeds under study. In addition Hanuš *et al.* (2011) observed that herd, year, and season showed a significant effect on

milk composition and its technological parameters (including RCT) of milk produced by dairy cows of the Czech Fleckvieh breed. Daviau *et al.* (2000) mentioned that shorter RCT was associated with a decrease in the content of protein, which usually associated also with a decrease in the content of casein. A significant effect of season and herd on rennet coagulation time and other parameters of milk technological quality in Holstein cows was observed also by Chládek *et al.* (2011).

The aim of this study was to determine the effect of barn airspace temperature on composition (i.e. contents of protein, fat, lactose and non fat solids) and technological parameters (titratable acidity, rennetability and curd quality) of bulk milk samples collected in herds of Czech Fleckvieh and Holstein breeds of cattle.

MATERIAL AND METHODS

The study was performed on two farms (A and B) in the South Moravian Region of the Czech Republic within the period from June 3rd 2010 to February 2nd 2011.

The herd on the first farm (A) consisted only from purebred Holstein (H) dairy cows (in average 350 head). In this herd, the average milk performance was 9,500 kg per lactation. The farm is situated in the village of Žabčice in a lowland area (GPS 49°0'51.786"N, 16°36'14.809"E) at the altitude of 179 m. All cows were kept together under identical conditions in a loose housing system with bedding and received a complete feeding ration *ad libitum*. Cows were milked twice a day at 4.00 and 16.00 h. This was the same barn as that used in experiments performed by Walterová *et al.* (2009).

On the other farm (B), only purebred dairy cows (in average 600 head) of the Czech Fleckvieh (CF) breed were raised. The average milk performance was 7,500 kg per lactation. The farm is situated in a lowland region in the village of Říčany, Moravia, Czech Republic (GPS 49°12'32.319"N, 16°23'42.666"E) in the altitude of 349 m. All animals were kept under identical conditions in a loose housing system with bedding and received also a complete feeding ration *ad libitum*. They were milked twice daily also at 4.00 and 16.00 h. This experiment took place in the same barn as that used by Erbez *et al.* (2010). On both farms were optimized diet according to Petrikovič and Sommer (2002). Feeding ration consisted from common used feeds in this region (corn silage, cereal meals, solvent oil meals, minerals and vitamins supplements).

Within a period of 36 weeks, bulk milk samples were collected in both herds once a week always on the same day. The samples represented a mixture of morning and evening milk. The average barn airspace temperature (BAT in °C) was measured on the day before milk sampling. Temperature measurements were performed every 15 minutes using three HOBO data loggers (H08-007-02, Onset Computer Corporation*), which were located

approx. 1.40m above the floor level in three different locations inside the barn to eliminate the effect of only one place of measuring.

On the next day, the average percentages of fat content (F), protein (P), lactate monohydrate (L), and solids non fat (SNF) were estimated in collected bulk milk samples together with values of titratable acidity (TA), rennetability (RCT), and curd quality (CQC). Milk rennetability was estimated using a „Nephelometric-turbidimetric test of milk coagulation“ (Chládek and Čejna, 2005). The test was performed using the preparation Laktochym 1:5000 (Milcom Tábor) in the dose of 1 ml per 50ml of milk (after the dilution of the renneting agent in the ratio 1:4). Curd quality (CQC) was evaluated after 60 minutes of incubation of 50 ml of renneted milk at 35 °C and compared with tabular values (Gajdůšek, 1999) using the scale from (1 = the best to 5 = the worst). TA was measured in a milk sample of 100 ml using an alkaline solution up to light pink colour of the mixture (in ml of the 0.25 molx1⁻¹ NaOHx100ml⁻¹). The method was performed pursuant provisions of the standard CSN 57 0530. Contents of P and F were estimated using the apparatus Milkoscope C5 (see the standard ČSN 57 0536).

For statistical analysis (by means of bi-factorial analysis of variance), programmes MS Excel and UNISTAT Version 5.1 were used.

The analyses carried on, including abbreviations and units of measurement were as follows:

| | |
|-----|---------------------------------------------|
| H | = Holstein |
| CF | = Czech Fleckvieh |
| P | = protein content (%), g.100g ⁻¹ |
| F | = fat content (%), g.100g ⁻¹ |
| L | = lactose (%), g.100g ⁻¹ |
| SNF | = solid non fat (%), g.100g ⁻¹ |
| BAT | = barn airspace temperature (°C) |
| RCT | = rennet coagulation time (in seconds) |
| CQC | = curd quality class |
| TA | = titratable acidity (°SH). |

RESULTS AND DISCUSSION

Values of mean, minimum, maximum and standard deviation (SD) of data from analysis of cow's milk composition, technological parameters and barn airspace temperature are shown in Table I. On both farms (n = 72), the average value of BAT was 12.08 °C and the standard deviation was 9.3 °C. On the farm A, the average BATs ranged from a minimum of -3.96 °C to the maximum of 28.51°C; for the whole period under study, the average value of BAT was 13.08 °C. On the farm B, the corresponding values of BATs ranged from -7.41 °C to +26.24 °C; for the whole period under study, the average value of BAT was 11.08 °C. This means that in some periods the monitored dairy cows were exposed to a heat stress (above all if BATs approached to the limit of 26 °C). Many authors (e.g. Berman *et al.*, 1985; Hahn, 1999 and West, 2003) reported that BATs above 23–26 °C were for dairy cattle critical and that caused a decrease in milk production. Some other authors, however, (e.g. Falta *et al.*, 2008; Vokřálová and Novák, 2005) demonstrated that for high-yielding dairy cows BATs of only 21 °C were critical and triggered the heat stress.

In both herds, the average values of P and its standard deviation were 3.46% and ± 0.20%, respectively. On the farm A, the average value of P was 3.33% (with the minimum and the maximum of 3.14% and 3.56%, respectively) while on the farm B it was 3.60% (with the minimum and the maximum of 3.33% and 3.83%, respectively). The difference between farms A and B was statistically highly significant (P < 0.01). The average values of F and its standard deviation for the whole period under study and both herds were 4.09% and ± 0.21%, respectively). On the farm A, the average F value was 4.06% (with the minimum and the maximum of 3.64% and 4.48%, respectively) while on the farm B it was 4.11% (with the minimum and the maximum of 3.67% and 4.41%, respectively). The difference between both farms was statistically non-significant (P > 0.05). From our observed values F

I: Mean, minimum, maximum and standard deviation of milk composition, technological properties and barn airspace temperature on both farms (A and B)

| Parameter | Total | | | | Farm A | | | Farm B | | | Signifi- cation |
|---------------------|-----------|-------|-------|-------|-----------|-------|-------|-----------|-------|-------|--------------------|
| | \bar{x} | SD | min. | max. | \bar{x} | min. | max. | \bar{x} | min. | max. | |
| RCT (second) | 211 | 15.69 | 160 | 240 | 213 | 185 | 240 | 209 | 160 | 240 | N.S. |
| CQC (class) | 1.53 | 0.50 | 1.00 | 2.00 | 1.53 | 1.00 | 2.00 | 1.53 | 1.00 | 2.00 | N.S. |
| TA (°SH) | 7.16 | 0.23 | 6.42 | 7.64 | 7.11 | 6.64 | 7.50 | 7.22 | 6.42 | 7.64 | N.S. |
| P (%) | 3.46 | 0.20 | 3.14 | 3.83 | 3.33 | 3.14 | 3.56 | 3.60 | 3.33 | 3.83 | ** |
| F (%) | 4.09 | 0.21 | 3.64 | 4.48 | 4.06 | 3.64 | 4.48 | 4.11 | 3.67 | 4.41 | N.S. |
| L (%) | 4.79 | 0.06 | 4.61 | 4.95 | 4.80 | 4.62 | 4.95 | 4.77 | 4.61 | 4.86 | N.S. |
| SNF (%) | 8.86 | 0.20 | 8.44 | 9.24 | 8.73 | 8.44 | 9.20 | 8.99 | 8.77 | 9.24 | ** |
| BAT (C°) | 12.08 | 9.30 | -7.41 | 28.51 | 13.8 | -3.96 | 28.51 | 11.08 | -7.41 | 26.24 | N.S. |

Signification: N.S. – non significant (P > 0.05); ** (P < 0.01)

RCT – rennet coagulation time; CQC – curd quality class; TA – titratable acidity; P – protein content; F – fat content; L – lactose content; SNF – solids non fat; BAT – barn airspace temperature, SD – standard deviation

and P, we can say that milk yield both breeds were higher than average in conditions of Czech Republic in comparison to data from the Milk Recording Scheme for the year 2010 according to Kvapilík *et al.* (2011). In the whole set of dairy cows and for the whole period under study, the average values of L and its standard deviation were 4.79% and $\pm 0.06\%$, respectively. On the farm A, the average value of L was 4.80% (with the minimum and the maximum of 4.62% and 4.95%, respectively). On the farm B, the corresponding value was 4.77% (with the minimum and the maximum of 4.61% and 4.86%, respectively). Also this difference between both farms was statistically non-significant ($P > 0.05$). In both herds and for the whole period under study, the average values of SNF and its standard deviation were 8.86% and $\pm 0.20\%$, respectively. On the farm A, the average value of SNF was 8.73% (with the minimum and the maximum of 8.44% and 9.02%, respectively). On the farm B, the corresponding value was 8.99% (with the minimum and the maximum of 8.77% and 9.24%, respectively). This difference between both farms statistically highly significant ($P < 0.01$).

For the whole period under study and both herds, the average values of RCT and its standard deviation were 211 sec and ± 15.69 sec. On farm A, the average value was 213 sec. (with the minimum and the maximum of 185 sec and 240 sec, respectively), while on the farm B it was 209 sec. (with the minimum and the maximum of 160 sec. and 240 sec, respectively). The difference between both farms was statistically non-significant ($P > 0.05$). As far as the values of CQC and its standard deviation for the whole study period were concerned, these were 1.53 ± 0.50 , respectively. The same average values were found out also in both herds and the minimum and the maximum values (i.e. Class 1 and Class 2, respectively) were recorded in both herds. On the farm A, the average value of TA was 7.11 °SH and ranged from 6.64 to 7.50 °SH; on the farm B, the corresponding values were 7.22 and 6.42–7.64 °SH, respectively. In both herds, the average values of TA and its standard deviation were 7.16 °SH and ± 0.23 , respectively. The difference between both farms was statistically non-significant ($P > 0.05$).


The correlation between milk content, technological parameters and barn airspace temperature on farms A and B are shown in Tab. II. These data indicate a marked effect of BAT on all parameters of milk composition and technological quality on both farms; non-significant was only the effect of BAT on L content on the farm A. RCT was negatively correlated with BAT on both farms ($r = -0.45$ and $r = -0.54$, respectively; $P < 0.01$). This means that the higher the value of BAT, the shorter that of RCT. Average values of summer BAT indicated that during this season, the limit of heat stress could be trespassed on some days (Falta *et al.*, 2008; Hanuš *et al.*, 2008). As mentioned by Daviau *et al.* (2008), the shorter RCT was associated with a decrease in P content and also of casein. It was found out in this study that lower values of RCT were associated with a lower content of protein above all in the summer season; however, our results do not correspond with data published by Jöudu *et al.* (2008); Ikonnen *et al.* (2004) and Sevi *et al.* (2001) who obtained opposite results. This could be partly explained on the base of high summer temperatures recorded in our study. This observation also corresponded with results published by Nájera *et al.* (2003). Regardless to differences existing between both farms, the value of CQC was positively correlated with BAT on both farms ($r = 0.42$ and $r = 0.36$ on farms A and B, respectively; $P < 0.01$) while that of TA was correlated negatively ($r = -0.39$ and $r = -0.44$ on farms A and B, respectively; $P < 0.01$). This effect of BAT on RTC on farms A and B is shown also in Fig. 1.

As far as the effect of BAT on values of P was concerned, the highest negative correlation coefficient was found out on both farms ($r = -0.86$ and $r = -0.88$ on farm A and B, respectively; $P < 0.01$). This effect of BAT on P content is obvious also in Fig. 2. Further, a negative coefficient of correlation was found out also between BAT and F content on both farms ($r = -0.77$ and $r = -0.87$ on farm A and B, respectively; $P < 0.01$). This effect of BAT on F content on farms A and B is illustrated also in Fig. 3. This trend in growth of P under conditions of decreasing temperatures was published by several authors (Hanus *et al.*, 2008; Dolejš *et al.*,


II: Correlation between milk content, technological properties and barn airspace temperature on farm A and farm B

| Farm A | | | | | | | | | |
|--------|-----|---------|---------|---------|---------|---------|---------|---------|---------|
| Farm B | I | RCT | CQC | TA | P | F | L | SNF | BAT |
| | RCT | 1 | -0.1215 | 0.1200 | 0.3627 | 0.5469 | -0.0199 | 0.2946 | -0.4478 |
| | CQC | -0.1858 | 1 | -0.2403 | -0.4292 | -0.3715 | -0.0670 | -0.4066 | 0.4198 |
| | TA | 0.0806 | -0.3225 | 1 | 0.3601 | -0.0047 | -0.3365 | 0.1844 | -0.3895 |
| | P | 0.6277 | -0.3922 | 0.5076 | 1 | 0.7107 | -0.0220 | 0.8877 | -0.8556 |
| | F | 0.5771 | -0.3301 | 0.3487 | 0.8584 | 1 | 0.0201 | 0.6249 | -0.7657 |
| | L | -0.4731 | 0.1691 | -0.2790 | -0.6701 | -0.5476 | 1 | 0.4028 | 0.0545 |
| | SNF | 0.6500 | -0.4191 | 0.4920 | 0.9566 | 0.8446 | -0.4672 | 1 | -0.7255 |
| | BAT | -0.5410 | 0.3599 | -0.4358 | -0.8824 | -0.8662 | 0.5699 | -0.8556 | 1 |

RCT – rennet coagulation time; CQC – quality of curd; TA – titratable acidity; P – protein content; F – fat content; L – lactose content; SNF – solids non fat; BAT – barn airspace temperature




1: Effect of barn airspace temperature on rennet coagulation time on farm A and B



2: Effect of barn airspace temperature on protein content on farm A and B

1996; Ng-Kwai-Hang *et al.*, 1984; Lacroix *et al.*, 1996). Kadzere *et al.* (2001) confirmed that during periods of warm weather, the percentage of milk protein decreased in all dairy cows. Moreover, McDowell *et al.* (1976) mentioned that if lactating dairy cows were transferred from a barn with air temperature of 18 to another with 30 °C, production of milk fat, solids non fat and milk protein decreased by 39.7; 18.9 and 16.9%, respectively. Ozrenk and Inci (2008) also observed that contents of protein and fat of milk change along the year and that the percent of

milk protein was positively correlated with that of milk fat. This observation was corroborated also in this study: it was found out that there was a positive correlation between contents of F and P (values of correlation coefficients on farms A and B were $r = 0.71$ and $r = 0.86$, respectively; $P < 0.01$). Further it was found out that there was a positive correlation between BAT and L content on farms B and A ($r = 0.05$; $P > 0.05$ and $r = 0.57$; $P < 0.01$, respectively). The result recorded on farm B differs from data published by Kadzere *et al.* (2001) who wrote that temperature



3: Effect of barn airspace temperature on fat content on farm A and B

did not affect the lactose percentage. This marked difference can be partly explained by the fact that the breed of cattle was different. On both farms, the correlation between BAT and SNF was also very high ($r = -0.73$ and $r = -0.86$ on farms A and B, respectively).

It can be therefore concluded that differences in RCT, as observed in our study (i.e. 13.5 %), were not the same as those recorded by Hanuš *et al.* (2010), respectively. Chládek *et al.* (2011) under different conditions and in different breeds (34.00%, resp. 22.7 %). The existence of significant differences among individual breeds support the opinion that the parameter „breed“ should be taken into account as one of factors that influence results of experiments focused on milk technological quality. An insignificant effect of breed on parameters of technological quality of milk observed in this study does not corresponds with data published by Hanuš *et al.* (2011) who recorded them in a study with a different breed of cattle.

It is obvious that similar trends in the growth and decrease of temperatures were observed in the course of this study on both farms. However, it can be concluded that average daily temperatures were nearly identical on both farms and that the average difference was approximately 2 °C. This

resulted above all from different localities, in which both farms were situated in the region of South Moravia. Thus, the differences in composition and technological parameters of milk resulted above all from different breeds: as compared with H dairy cows, those of CF breed produced statistically significantly ($P < 0.01$) higher percentages of P (by +0.27 %) and SNF (by +0.26 %). The other differences were statistically non-significant ($P > 0.05$): BAT (−2 °C), RCT (−4 sec.), CQC (0), TA (+0.11 °SH), F (+0.05 %) and L (−0.03 %). As far as the effect of temperature is concerned, the calculated values of correlation coefficients (and also the plotted graphs) indicate a marked effect of BAT on RCT, F and P. At lower temperatures, this effect was always markedly negative (i.e. higher in case of F, P and longer in case of RCT) while at high temperatures it was less pronounced (i.e. lower and shorter). It is also necessary to remember that, within the period under study, milk composition and its technological parameters were markedly influenced by the temperature (BAT), which could further deepen differences existing between individual breeds. Regardless to differences in average values of parameters under study, these trends were similar in both breeds.

SUMMARY

This study was performed in two herds reared on two farms in the territory of South Moravian Region (Czech Republic). On farm A, there was a herd of Holstein cattle while on farm B the dairy cows were on CF breed. Altogether 72 samples of bulk milk were collected within the period beginning on 3 July 2010 and ending on 2 February 2011. Averages values of technological milk quality parameters were as follows: rennet coagulation time (RCT) 211 sec., titratable acidity (TA) 7.16 °SH, quality of curd (CQC)

1.53; fat content (F) 4.09%, protein content (P) 3.46%, lactose carbohydrate content (L) 4.79% and solids non fat content (SNF) 8.86%. The average daily temperature was 12.08 °C. The effect of season on nearly all parameters under study was significant (the only exception was L content on farm A). When comparing the results obtained during the year, the significantly shortest RCT value was recorded in summer while in winter it was the longest. In summer, values of F, P and SNF were the lowest. In the autumn, the lowest values of L were recorded while those of F and P were the highest. In winter, the values of SNF were the highest at all. The effect of breed on nearly all parameters under study was non-significant the only exception represented contents of P and SNF). Maximum differences between both breeds were as follows: RCT 80 sec, TA 1.22 °SH, QC 1, F 0.84 g.100g⁻¹, P 0.69 g.100g⁻¹, L 0.34 g.100g⁻¹, SNF 0.8 g.100g⁻¹. When comparing the course of temperatures in both localities, it is possible to conclude that trends in growth and decrease of temperatures was nearly the same during the study period and that it was equal to 2 °C. This resulted above all from different localities, in which both farms were situated in the region of South Moravia. It is also possible to concluded that the shortest values of RCT were recorded on both farms during the summer season.

Acknowledgements

This study was supported by the Research Project No. MSM6215648905 „Biological and technological aspects of sustainability of controlled ecosystems and their adaptability to climate change“, which was financed by the Ministry of Education, Youth and Sports (MŠMT) of the Czech Republic and by the project MŠMT Kontakt ME 09081.

REFERENCES

- BEATTY, D. T., BARNES, A., TAYLOR, E., PETHICK, D., MCCARTHY, M., MALONEY, S. K., 2006: Physiological response of *Bos taurus indicus* cattle to prolonged, continuous heat and humidity. *Journal of Animal Science*, 84: 972–985. ISSN 0021-8812.
- BERMAN, A., FOLMAN, Y., KAIM, M., MAMEN, M., HERZ, Z., WOLFENSON, D., ARIELI, A., GRABER, Y., 1985: Upper critical temperatures and forced ventilation effects for high-yielding dairy cows in a subtropical climate. *Journal of Dairy Science*, 68: 1488–1495. ISSN 0022-0302.
- CASATI, M. R., CAPPALÀ, V., CALAMARI, L., CALEGARI, F., FOLLI, G., 1998: Effects of the season on milk yield and some milk characteristics in cows. *Scienza e Tecnica Lattiero-casearia*, 49: 7–25. ISBN 88-464-7408-2.
- ČSN 570530, 1972: Methods for testing of milk and dairy products. Standart ČNI prague, (in Czech).
- ČSN 570536, 1999: Determination of milk composition by mid-infrared analyzer. (In Czech).
- DAVIAU, C., FAMELART, M. H., PIERRE, A., GOUDÉDRANCHE, H., MAUBOIS, J. L., 2000: Rennet coagulation of skim milk and curd drainage: Effect of pH, casein concentration, ionic strength and heat treatment. *Lait* 80, 4: 397–415. ISSN 1958-5586.
- DAVIS, M. S., MADER, T. L., HOLT, S. M., PARKHUST, A. M., 2003: Strategies to reduce feedlot cattle heat stress: Effects on tympanic temperature. *Journal of Animal Science*, 81: 649–661. ISSN 0021-8812.
- De MARCHI, M., DAL ZOTTO, R., CASSANDRO, M., BITTANTE, G., 2007: Milk coagulation ability of five dairy cattle breeds. *Journal of Dairy Science*, 90, 8: 3986–3992. ISSN 0022-0302.
- DOLEJŠ, J., TOUFAR, O., KNÍŽEK, J., 1996: The temperature of rearing environment affects on milk chemical composition. (In Czech), *Náš Chov*, 7: 20. ISSN 0027-8068.
- ERBEZ, M., FALTA, D., CHLÁDEK, G., 2010: The relationship between temperature and humidity outside and inside the permanently open-sided cow's barn. *Acta Universitatis agriculturae et silviculturae Mendelianae Brunensis*. LVIII, 5: 91–96. ISSN 1211-8516.
- FALTA, D., WALTEROVÁ, L., SKÝPALA, M., CHLÁDEK, G., 2008: Effect of stable microclimate on milk production of Holstein cows on the 2nd and 3rd lactation. *Animal welfare, etológia és tartástechnológia*. [online] 7, URL: <http://www.animalwelfare.szie.hu/Cikkek/200802/Szarvasmarha/AWETH2008104110.pdf>.
- GADER, A. Z., AHMED, M. K. A., MUSA, L. M. A., PETERS, K. J., 2007: Milk yield and reproductive performance of Friesian cows under Sudan tropical conditions. *Archiv Tierzucht*. 50, 155–64. ISSN 0003-9438.
- GAJDŮŠEK, S., 1999: Milk processing II – practise. Mendel University in Brno, (in Czech), p. 92.
- HAENLEIN, G., 2003: Nutritional value of dairy products of ewe and goat milk. Retrieved January 28, from <http://ag.udel.edu/extension/information/goatmgt/gm-10.htm>.
- HAHN, G. L., 1999: Dynamic response of cattle to thermal heat loads. *Journal of Animal Science*, 77: 10–20. ISSN 0021-8812.
- HANUŠ, O., FRELICH, J., JANŮ, L., MACEK, A., ZAJÍČKOVÁ, I., GENČUROVÁ, V., JEDELSKÁ,

- R., 2007: Impact of different milk yields of cows on milk quality in Bohemian spotted cattle. *Acta Veterinaria Brno*, 76: 563–571. ISSN 1801-7576.
- HANUŠ, O., FRELICH, J., TOMÁŠKA, M., VYLETĚLOVÁ, M., GENČUROVÁ, V., KUČERA, J., TRINÁCTÝ, J., 2010: The analysis of relationships between chemical composition, physical, technological and health indicators and freezing point in raw cow milk. *Czech Journal of Animal Science*, 55, 1: 11–29. ISSN 1212-1819.
- HANUŠ, O., KUČERA, J., YONG, T., CHLÁDEK, G., HOLÁSEK, R., TRINÁCTÝ, J., GENČUROVÁ, V., SOJKOVÁ, K., 2011: Effects of sires on wide scale of milk indicators if first calving Czech Fleckvieh cows. *Archiv Tierzucht*, 54, 1: 36–50. ISSN 0003-9438.
- HANUŠ, O., VYLETĚLOVÁ, M., GENČUROVÁ, V., JEDELSKÁ, R., KOPECKÝ, J., NEZVAL, O., 2008: Hot stress of Holstein dairy cows as substantiv factor of milk composition. *Scientia Agriculturae Bohemica*, 39 (4): 310–317. ISSN 0582-2343.
- CHLÁDEK, G., ČEJNA, V., 2005: Measuring of rennet coagulation time by Nephelo-turbidimetric sensor of milk coagulation. *Mléko a sýry 2005*, (In Czech), Prague, 127–130. ISBN 80-8623-8-48-2.
- CHLÁDEK, G., ČEJNA, FALTA, D., MÁCHAL., 2011: Effect of season and herd on rennet coagulation time and other parametres of milk technological quality in Holstein dairy cows. *Acta Universitatis agriculturae et silviculturae Mendelianae Brunensis*, LIX, 5: 113–118. ISSN 1211-8516.
- IKONNEN, T., MORRI, S., TYRISEVÁ, A. M., ROUTTINEN, O., OJALA, M., 2004: Genetic and phenotypic correlations between milk coagulation properties, milk production traits, somatic cell count, casein content and pH of milk. *Journal of Dairy Science*, 87: 458–467. ISSN 0022-0302.
- JANŮ, L., HANUŠ, O., FRELICH, J., MACEK, A., ZAJÍČKOVÁ, I., GENČUROVÁ, V., JEDELSKÁ, R., 2007: Influence of different milk yields of Holstein cows on milk quality indicators in the Czech Republic. *Acta Veterinaria Brno*, 76, 4: 553–561. ISSN 1801-7576.
- JÖUDU, I., HENNO, M., KAART, T., PÜSSA, T., KÄRT, O., 2008: The effect of milk protein on the rennet coagulation properties of milk from individual dairy cows. *International Dairy Journal*, 18, 9: 964–967. ISSN 0958-6946.
- KADZERE, C. T., MURPHY, M. R., SILANIKOVE, N., MALTZ, E., 2001: Heat stress in lactating dairy cows: a review. *Livestock Production Science*, 77: 59–91. ISSN: 0301-6226.
- KVAPILÍK, J., RŮŽIČKA, Z., BUCEK, P., 2011: Ročenka. Chov skotu v České republice. 36–37. ISBN 978-80-904131-6-0.
- LACROIX, C., VERRET, P., PAQUIN, P., 1996: Regional and seasonal variations of nitrogen fraction in commingled milk. *International Dairy Journal*, 6: 947–961. ISSN: 0958-6946.
- MADER, T. L., DAVIS, M. S., 2004: Effect of management strategies on reducing heat stress of feedlot cattle: Feed water intake. *Journal of Animal Science*, 82: 3077–3087. ISSN 0021-8812.
- MAUST, L. E., MCDOWELL, R. E., HOOVEN, N. W., 1972: Effects of summer weather on performance of Holstein cows in three stages of lactation. *Journal of Dairy Science*, 55: 1133–1139. ISSN 0022-0302.
- MCDOWELL, R. E., HOOVEN, N. W., CAMOENS, J. K., 1976: Effects of climate on performance of Holstein in first lactation. *Journal of Dairy Science*, 59: 965–973. ISSN 0022-0302.
- MITLÖHNER, F. M., GALYEAN, M. L., MCLOGN, J. J., 2002: Shade effects on performance, carcass traits, physiology, and behavior of heat-stressed feedlot heifers. *Journal of Animal Science*, 80: 2043–2050. ISSN 0021-8812.
- NÁJERA, A. I., DE RENOBABLES, M., BARRON, L. J. R., 2003: Effects of pH, temperature, CaCl₂ and enzyme concentrations on the rennet-clotting properte of milk: a multifactorial study. *Food Chemistry*, 80, 3: 345–352. ISSN 0308-8146.
- NG-KWAI-HANG, K. F., HAYES, J. F., MOXLEY, J. E., MONARDES, H. G., 1984: Variability of test day milk production and composition and relation of somatic cell counts with yield and compositional ganges of bovine milk. *Journal of Dairy Science*, 67: 361–366. ISSN 0022-0302.
- NOVÁK, P., VOKŘÁLKOVÁ, J., BROUČEK, J., 2009: Effects of the stage and number of lactation on milk yield of dairy cows kept in open barn during high temperatures in summer months. *Archiv Tierzucht*, 52, 6: 574–586. ISSN 0003-9438.
- OZRENK, E., INCI, S. S., 2008: The effect of seasonal variation on the composition of cow milk in Van Province. *Pakistan Journal of Nutrition*, 7 (1): 161–164. ISSN 1680-5194.
- PETRIKOVIČ, P., SOMMER, A., 2002: Potreba živín pre hovézí dobytok. *Výskumný ústav živočišnej výroby Nitra*, 6: 3–60. ISBN 80-88872-21-9.
- SEKERDEN, O., 1999: Effects of calving season and lactation order on milk yield and milk components in Simmental cows. *Turkish Journal of Veterinary and Animal Sciences*, 23: 79–86. ISSN 1300-0128.
- SEVI, A., ANNICCHIARICO, G., ALBENZIO, M., TAIBI, L., MUSCIO, A., DELL'AQUILA, S., 2001: Effects of solar radiation and feeding time on behavior, immune responses and production of lactating ewes under high ambient temperature. *Journal of Dairy Science*, 84: 629–640. ISSN 0022-0302.
- TUCKER, H. A., 1989: Photoperiod affects intake, growth and milk production of cattle. *Feedstuffs*, 61: 15–16.
- VOKŘÁLKOVÁ, J., NOVÁK, P., 2005: Climatic extrens and lactation. (In Czech), *Náš chov*, 9: 40–42. ISSN 0027-8068.
- WALTEROVÁ, L., ŠAROVSKÁ, L., FALTA, D., CHLÁDEK, G., 2009: Relation between some climate parametres inside and outside the stable in the course of the year. *Acta Universitatis agriculturae et silviculturae Mendelianae Brunensis*, LVII, 4: 125–132. ISSN 0165-7380.

- WEST, J. W., 2003: Effects of Heat-Stress on production in Dairy Cattle. *Journal of Dairy Sciences*, 86: 2131–2144. ISSN 0022-0302.
- YETISMEYEN, A., 2000: Milk technology. Ankara, Ankara University Agriculture Faculty Press No: 1511.

Address

Ing. Ondřej Polák, Ing. Daniel Falta, Ph.D., prof. Ing. Gustav Chládek, CSc., Ústav chovu a šlechtění zvířat, Mendelova univerzita v Brně, Zemědělská 1, 613 00 Brno, Česká republika, doc. Ing. Oto Hanuš, Ph.D., Výzkumný ústav pro chov skotu Rapotín, Výzkumníků 267, 788 13 Víkřovice, Česká republika, e-mail: xpolak@mendelu.cz

