

OCCURRENCE OF MASTITIS PATHOGENS IN RELATION TO SOMATIC CELLS

Marcela Vyletřlová Klimeřová, Oto Hanuš, Lucie Hasoňová, Petr Roubal,
Ivan Manga, Ludmila Nejšlechbová

Received: March 27, 2013

Abstract

VYLETŘLOVÁ KLIMEŘOVÁ MARCELA, HANUŠ OTO, HASOŇOVÁ LUCIE, ROUBAL PETR, MANGA IVAN, NEJŠLECHBOVÁ LUDMILA: *Occurrence of mastitis pathogens in relation to somatic cells*. Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis, 2013, LXI, No. 5, pp. 1505–1511

There were examined 161 cows from 4 farms in total. The suspect animals were selected according to viscosity test results, clinical symptoms and somatic cell count (SCC). Milk samples were examined for the presence of pathogens and for SCC. 55 mastitis pathogens were identified. The most frequently isolated species was *Enterococcus faecalis* ($n = 20$), followed by *Staphylococcus aureus* ($n = 6$) and *Streptococcus uberis* ($n = 5$). The SCC ranged from 9 to 24 204 ths.ml⁻¹. There was positive occurrence of bacteria genus *Staphylococcus* and *Enterococcus* at lower SCC (50 ths.ml⁻¹) and at higher SCC numbers (> 300 ths.ml⁻¹) bacteria genus *Streptococcus*, *Enterobacter* and *Escherichia coli*. Differences in SCC were significant ($P < 0.001$) in negative samples xg 131 SCC versus 491 for positive, 611 for staphylococci and 464 ths.ml⁻¹ for other positive. SCC discrimination limit for practical likelihood of pathogen occurrence estimation in infectious sample groups was calculated. This limit for suspicion of infection is 159 for positive group, 113 for staphylococci and 174 ths.ml⁻¹ for other positive. This could be possible to recommend the value 174 ths.ml⁻¹ for practical use with target to apply preventive or curative measures.

cow's milk, *Staphylococcus aureus*, *Staphylococcus haemolyticus*, *Streptococcus uberis*, discrimination limit

The incidence of mastitis pathogens is a persistent problem not only in dairy cows but also for other dairy animals. For breeders it always presents the economic losses. Rajala-Schultz *et al.* (1999) studied the effect of mastitis disease on milk yields and found that the daily loss during the first 2 wk after the occurrence of mastitis varied from 1.0 to 2.5 kg, and the total loss over the entire lactation varied from 110 to 552 kg. The milking hygiene control, which is directly associated with the development disease caused by environmental microorganisms, often sticking mainly to time and financial cost of laboratory tests. The higher incidence of environmental microorganisms, which include *Staphylococcus aureus* and *Streptococcus uberis*, can be explained by the development and change of milking technology, when the pipeline milking equipment in stall occurred more frequent contamination of the udder by hands of nursing

staff or by handling with milking equipment. Therefore, the milking equipment influences whole row of indicators by its function and use, for instance by overmilking (Pařilová *et al.*, 2010, 2011), and can be factor and vector of mastitis origination. Aggravated function grows up mastitis occurrence and subsequently also somatic cell count (Hanusř and Ticháček, 1997). The current types of milking parlours provide better comfort and hygiene of milking (Vyletřlová *et al.*, 2009; Janřtová *et al.*, 2011). Beside good quality of milking equipment function also other ways of reduction of mastitis occurrence are looking for as cow vaccination for instance (Touřová *et al.*, 2011).

One of the respected indicators of health status of the mammary gland (Sawa and Piwczynski, 2002; Berry *et al.*, 2006; Heck *et al.*, 2009) is the somatic cells count (SCC) similarly as mammary associated isotype of serum amyloid A (Kováč *et al.*, 2011).

Golebiewski *et al.* (2011) found the higher average SCC value in long-term observation for Polish Holstein-Friesian as compared to Montbéliarde breed (642 and 455 ths.ml⁻¹). Similar conclusion published Frelich and Šlachta (2011) for Holstein and Czech Fleckvieh without interaction to farm and seasonal effect. The permissible SCC limit value in a bulk milk sample is < 400 thousands in 1 ml of raw cow's milk. As a healthy mammary gland is generally taken to mean that, where the SCC is around 100 ths.ml⁻¹ in quarter milk sample. Sheldrake *et al.* (1983) determined the number of somatic cells in uninfected mammary gland from 83 ths.ml⁻¹ (from the 35th day after calving) to 160 ths.ml⁻¹ (285th day). The higher values indicate a possible infection of the mammary gland. Reneau (1986) in his work provides the value of SCC 283 ths.ml⁻¹ as a limit value suitable for the determination of suspicion from subclinical mastitis occurrence. The less numbers of SCC (250 and 228 ths.ml⁻¹) indicate also Andrews *et al.* (1983) and Dohoo *et al.* (1981) as the threshold for selection in the case of treatment of mastitis in cows. The similar relationship (but in the case of the bulk milk samples) describe in their work Benda *et al.* (1997). They estimated 1.7% mastitis diseases incidence caused by *Staphylococcus aureus* at the SCC 160 ths.ml⁻¹, and 43.5% incidence at the SCC 410 ths.ml⁻¹. In the case of streptococcal mastitis caused by *Streptococcus agalactiae* estimated 1% incidence of mastitis at SCC 160 ths.ml⁻¹, and 24.6% at SCC 400 ths.ml⁻¹.

In time of areal application of antibiotic mastitis therapy in dairy cows and increase of pathogen strain resistance against these medicaments the topicality of return to selective cure and also importance of methods which support this kind of therapy is growing up. At present, much attention is given to methicillin-resistant staphylococci strains, especially *S. aureus* (MRSA), which represent a serious problem in human and veterinary medicine (Holmes and Zadoks, 2011). These strains, which are associated with livestock, are called as LA-MRSA. The spread of resistant strains in cattle can be problematic in the case of farms with breeding of dairy and meat animals. Because the transfer of resistance going horizontally, the spread of resistant strains is also possible through working staff (Holmes and Zadoks, 2011). An improvement of methods for support of selective cure could be effective in terms of cost saving and mentioned risk reduction as well.

Aim of this work was to assess the relationship between mammary gland pathogen occurrence and somatic cell count (SCC) and possibility to find SCC discrimination limit for estimation of start for subclinical mastitis treatment according to SCC under current conditions.

MATERIAL AND METHODS

Animals and milk sample collection

The suspect animals were selected by responsible worker of dairy production and according to NK test (viscosity) results, clinical symptoms and SCC. There were mainly the higher lactation cows (> 1st lactation) with subclinical mastitis. Milk samples were collected from all four teats into sterile sample containers, kept in a cooler at 4 °C, transported to the laboratory and then immediately processed. There were examined 161 cows from four farms in total. Holstein and Czech Fleckvieh dairy cows were included in the experiment. The mean herd milk yield varied from 5 600 to 8 900 kg per standard lactation. The binding stabling with pipeline milking was used in two stables and free stabling with milking parlour in other two stables. All animals were milked twice a day.

Microbiological laboratory analyses

For microbiological analysis of samples there were used Blood agar, Edwards and Endo agar (Oxoid, Basingstoke, UK). The cultivation of the samples proceeded at 36 °C for 24 hours. Suspicious colonies were further isolate on Blood agar and cultivated at 36 °C for 24 hours. For species identification there were used biochemical tests STREPTOtest, STAPHYtest and ENTEROtest and identification program TNWpro7.0 (Erba Lachema, CZ). In addition, in strains confirmed as *S. aureus* was detected *mecA* gene, encoding resistance to methicillin, using PCR method (Boşgelmez-Tinaz *et al.*, 2006). The determination of SCC in individual milk samples was performed according to CSN EN ISO 13366-2 (2007) using the Somacount instrument (Bentley Chasca Minnesota, USA).

Statistical methods

The SCC classification into classes according to the positive and negative findings of microbiological pathogens was used for statistical evaluation of the results. The classes were created by respectable, literature-based professional practice. Somatic cell counts were evaluated in their original values (thousands ml⁻¹), but also in logarithmically transformed form. This was done for a legitimate presumption absence of normal frequency distribution of SCC results in sets of individual milk samples (Ali and Shook, 1980; Raubertas and Shook, 1982; Reneau *et al.*, 1983, 1988; Reneau, 1986; Wiggans and Shook, 1987). For the same reason, for statistical hypothesis testing of differences between groups of samples (t-test on log SCC), there were used geometric means and medians (in the position of the mean values) because of the given purpose. These are more representative characteristics than a simple arithmetic indicator.

RESULTS AND DISCUSSION

Pathogen species identification

Tab. I shows the results of species identification of the mastitis pathogens. We have confirmed 52 positive dairy cows from all of 161 tested animals and identified 55 mastitis pathogens. The most frequently isolated species was *Enterococcus faecalis* ($n = 20$), followed by *Staphylococcus aureus* ($n = 6$) and *Streptococcus uberis* ($n = 5$). In *S. aureus* strains, the presence of the *mecA* gene, or genes for enterotoxins production was not found. A similar incidence of mastitis pathogens and negative incidence of MRSA was confirmed also in previous work (Vyleťelová *et al.*, 2011).

The interesting thing is the finding of only one case of pathogen *Streptococcus agalactiae* from all of positive samples. Previously this pathogen was the main cause of contagious mastitis with high SCC together with *S. aureus* (Erskine *et al.*, 1987; Benda *et al.*, 1997; Keefe, 1997). Today is *S. agalactiae* after changing technologies, housing and milking hygiene regimes, essentially replaced by other environmental pathogen *S. uberis*. Next to *S. aureus* and *S. uberis* are the most important other species of pathogenic microorganisms such as *E. coli*, coagulase-negative staphylococci CNS, *Corynebacterium bovis* (Bradley, 2002; Pitkälä *et al.*, 2004; Vyleťelová, 2006; Vasil, 2009; Kalmus *et al.*, 2011).

Occurrence of mastitis pathogens in relation to somatic cells

Somatic cell count ranged from 9 to 24204 thousands (ths.) in 1 ml. In Table I, there is divided incidence of mastitis pathogens in relation to somatic cells into three groups: I) positive / negative mastitis incidence of disease in the case of the $\text{SCC} \leq 100 \text{ ths.ml}^{-1}$; II) incidence of mastitis disease if $\text{SCC} > 100$ and $\leq 283 \text{ ths.ml}^{-1}$; III) incidence of mastitis disease in case of $\text{SCC} > 283 \text{ ths.ml}^{-1}$. As regards, the positive or negative mastitis disease incidence in the case of I. and II. classes, significant differences are apparent. This also confirms the mentioned limit value for the determination of suspicion of subclinical mastitis (Reneau, 1986). However, differences are between species proportion of isolated bacterial strains.

From the results, it is interesting that at lower SCC (about 50 ths.ml^{-1}) were present bacteria of the genus *Staphylococcus*, followed by bacteria of genus *Enterococcus* (75 ths.ml^{-1}), and only at higher numbers of SCC (346, 554 and 594 ths.ml^{-1} respectively) bacteria genus *Streptococcus*, *Enterobacter* and *Escherichia coli*.

From all results it was found 23 negative results (14%) in samples with higher SCC than 400 ths.ml^{-1} (from 430 to 9006 ths.ml^{-1}). The absence of pathogens in this case may be caused by another load of mammary glands (stress, nutrition, milking equipment). Similar experiences, but with quarter (individual) samples describe Emanuelson and

Wever (1989), when the SCC correctly classified 82.9% of all cases of suspected bacterial infection.

Results of SCC statistical analyses

Statistical analysis results of SCC sets are shown in tables (Tabs. II and III) and in graph (Figure 1). In Tab. II, there are shown mean SCC values for individual classes of milk samples from health and infected mammary glands and in Table III significance of differences between them. It is remarkable at typically higher arithmetic means of SCC in individual infected classes (from 1923 to 3728 ths.ml^{-1}) and in contrast to non infected samples (295 ths.ml^{-1}) that their relative variability is quite balanced (v_x from 163.9 to 234.8 versus 173.5%). Also values of geometrical means (Tab. II) and medians (Fig. 1) of SCC individual classes are characteristically very well-balanced compared to values of arithmetical means. Therefore, these values are more representative, more reliable and hence preferable for interpretation. This conclusion is in accordance with previous results and estimations by Ali and Shook, (1980), Raubertas and Shook (1982), Reneau *et al.* (1983, 1988), Reneau (1986) and Wiggins and Shook (1987).

Differences in SCC (arithmetic and geometric means) according to pathogen classes (Tab. II) were statistically significant ($P < 0.001$) first of all as for about differences non infection group against infected groups which are represented by pathogens (Tab. III; x_g 131 SCC versus 491 for positive, 611 for staphylococci and 464 ths.ml^{-1} for other positive). These significant differences in SCC are as group median with negative findings against group medians with positive findings shown in Figure 1. Insignificant difference ($P > 0.05$; Tab. II, III and Fig. 1) in SCC was noted only between infected groups with positive pathogen findings (staphylococci versus other pathogens, x_g 611 and 464 ths.ml^{-1}).

In the file, there was minority representation of *S. agalactiae* infection, which marked again highest SCC loading ($15\,199 \text{ ths.ml}^{-1}$; Table I). Under mentioned conditions which are again in good accordance with current general proportions of mastitis ethiology (Vyleťelová, 2006; Vasil, 2009) the staphylococci infection means higher milk loading by somatic cells (x_g 611 ths.ml^{-1} ; Tab. II) and its quality aggravation than in the case of other infection (x_g 464 ths.ml^{-1} ; Tab. II). Of course, this fact increases economical relevancy of *S. aureus* infection.

It is possible to calculate a SCC discrimination limit (on 95% conventional level of probability) for practical likelihood of pathogen occurrence estimation in infectious sample groups under presumption of normal frequency distribution of logarithmic transformed SCC values (comparable obliqueness and acuteness of data files). This SCC discrimination limit for suspicion of infection is 159 ths.ml^{-1} for positive group (Tab. II) according to procedure: $\text{negative } 2.1157 + 0.5354 \times 1.96 = 3.165084$; $\text{positive } 2.6911 - 0.7407 \times 1.96 = 1.239328$; 3.165084

– $1.239328 = 1.925756/2 = 0.962878 + 1.239328 = 2.202206 = 159 \text{ ths.ml}^{-1}$ of SCC. Along the same procedure for sample groups staphylococci and other positive these limits are 113 and 174 ths.ml^{-1} . It is evident that these limits are mutually relatively similar. Perhaps this is possible to recommend the value 174 ths.ml^{-1} as limit for suspicion of infection and for practical use with target to apply selective mastitis antibiotic treatment or other preventive or curative measures. This could be used for instance as indication for selective antibiotic dry cow therapy in case of its exceeding as compared to SCC lactation geometric mean. This limit is lower as compare to limit 283 ths.ml^{-1} which was defined previously in dependence on milk yield losses by Reneau *et al.* (1983 and 1988), Reneau (1986) and Wiggans and Shook (1987) for similar purposes. However, this estimation is more similar to limits which were previously mentioned by Dohoo *et al.* (1981), Sheldrake *et al.* (1983), Andrews *et al.* (1983) and also Benda *et al.* (1997). One of the reasons for this could be also the development (changes) in pathogen and mastitis situation in cow herds (Sawa and Piwczynski, 2002; Berry *et al.*, 2006; Vyleťelová, 2006; Heck *et al.*, 2009; Vasil, 2009; Frelich and Šlachta, 2011; Golebiewski *et al.*, 2011) during included years in comparison.

CONCLUSIONS

From these obtained results and in consequence of related results by Vyleťelová *et al.* (2009 and 2011) there is possible to conclude as follows:

- *Staphylococcus aureus* is permanent risk in veterinary medicine and its systematic checking should be important part of HACCP and in milk primary production;
- during last years it is clear from results of development of mastitis pathogens that *Staphylococcus aureus* is henceforth one of main causes of mastitis and *S. uberis*, *E. faecalis* and CNS happen important microorganisms in mastitis pathogenesis;
- the highest frequency of mammary gland disorder was in subclinical mastitis and *Staphylococcus aureus* was the most frequently identified pathogen. *Streptococcus uberis*, *Staphylococcus haemolyticus* and *Streptococcus agalactiae* were following pathogens in order according to their frequency of occurrence.

In general the results pointed out by good correspondence of their main characteristics the interpretation importance of milk sample microbiological investigation on pathogen occurrence in indicated cases for possibility how to put into effect the preventive and curative measures in mastitis elimination including the investigation of pathogen antibiotic sensitivity. This is undervalued relatively and improperly by farmer practice today. In the fact it means change of present routine approach.

SUMMARY

The aim of this work was to assess the relationship between mammary gland pathogen occurrence and somatic cell count (SCC) and possibility to find SCC discrimination limit for estimation of start for subclinical mastitis treatment according to SCC under current conditions. The suspect animals were selected according to NK test (viscosity) results, clinical symptoms and SCC. There were mainly the higher lactation cows (> 1st lactation) with subclinical mastitis. Milk samples were collected from all four teats and there were examined 161 cows from four farms in total. Holstein and Czech Fleckvieh dairy cows were included in the experiment. For microbiological analysis there were used standard cultivation agars and the isolated species were identified using biochemical tests STREPTOtest, STAPHYtest and ENTEROtest and identification program TNWpro7.0. In addition, in strains confirmed as *S. aureus* was detected *mecA* gene, encoding resistance to methicillin, using PCR method. The determination of SCC in individual milk samples was performed using the Somacount instrument. The SCC classification into classes according to the positive and negative findings of microbial pathogens was used for statistical evaluation of the results. For the same reason, for statistical hypothesis testing of differences between groups of samples (t-test on log SCC), there were used geometric means and medians (in the position of the mean values) because of the given purpose. We have confirmed 52 positive dairy cows and identified 55 mastitis pathogens from our results. The most frequently isolated species was *Enterococcus faecalis* (n = 20), followed by *Staphylococcus aureus* (n = 6) and *Streptococcus uberis* (n = 5). In *S. aureus* strains, the presence of the *mecA* gene, or genes for enterotoxins production was not found. Somatic cell count ranged from 9 to 24 204 ths. in 1 ml. We have divided incidence of mastitis pathogens in relation to somatic cells into three groups: I) positive / negative mastitis incidence of disease in the case of the $\text{SCC} \leq 100 \text{ ths.ml}^{-1}$; II) incidence of mastitis disease if $\text{SCC} > 100$ and $\leq 283 \text{ ths.ml}^{-1}$; III) incidence of mastitis disease in case of $\text{SCC} > 283 \text{ ths.ml}^{-1}$. As regards, the positive or negative mastitis disease incidence in the case of I. and II. classes, significant differences are apparent. From the results, it is interesting that at lower SCC (50 ths.ml^{-1}) were present bacteria of the genus *Staphylococcus*, followed by bacteria of genus *Enterococcus* (75 ths.ml^{-1}), and only at higher numbers of SCC (346, 554 and 594 ths.ml^{-1} respectively) bacteria genus *Streptococcus*, *Enterobacter* and

Escherichia coli. From all results it was found 23 negative results (14%) in samples with higher SCC than 400 ths.ml⁻¹ (from 430 to 9 006 ths.ml⁻¹). The absence of pathogens in this case may be caused by another load of mammary glands (stress, nutrition, milking equipment). Differences in SCC according to pathogen classes were significant ($P < 0.001$; negative xg 131 SCC versus 491 for positive, 611 for staphylococci and 464 ths.ml⁻¹ for other positive). Insignificant difference in SCC was noted only between infected groups with positive pathogen findings (staphylococci versus other pathogens, xg 611 and 464 ths.ml⁻¹). It increases economical relevancy of *S. aureus* infection. It is possible to calculate a SCC discrimination limit for practical likelihood of pathogen occurrence estimation in infectious

I: Results of genus and species identification of mastitis pathogens, the lowest somatic cell count (SCC in ths.ml⁻¹) at pathogen confirmation

finding/SCC	I. ≤ 100	II. $> 100 \leq 283$	III. > 283	SCC ths.ml ⁻¹
negative (interval)	47	36	26	9–3 379
positive (interval)	6	11	38	52–24 204
<i>Staphylococcus aureus</i>	-	2	4	216
<i>Staphylococcus haemolyticus</i>	1	-	1	56
<i>Staphylococcus simulans</i>	1	-	-	64
<i>Staphylococcus xylosus</i>	2	-	-	52
<i>Staphylococcus intermedius</i>	-	-	1	711
<i>Enterococcus faecalis</i>	-	7	13	104
<i>Enterococcus faecium</i>	-	1	2	187
<i>Enterococcus gallinarum</i>	1	-	1	87
<i>Enterobacter amnigenus</i> bv. 1	-	1	-	123
<i>Escherichia coli</i>	-	-	1	594
<i>Streptococcus uberis</i>	-	-	5	594
<i>Streptococcus agalactiae</i>	-	-	1	15 199
<i>Staphylococcus</i> sp.	-	-	1	4 165
<i>Streptococcus</i> sp.	-	-	7	346
<i>Enterobacter</i> sp.	-	-	1	554
<i>Enterococcus</i> sp.	1	-	-	75

II: Statistical characteristics of case groups of somatic cell counts (SCC in ths.ml⁻¹) in milk with various mastitis state of mammary gland of animals with different pathogen occurrence respectively


Parameter	total	negative	positive	staphylococci	other positive
n	161	97	64	13	51
x	1 088	295	2 290	3 728	1 923
sd	3 241	511	4 882	6 109	4 517
vx	297.8	173.5	213.2	163.9	234.8
x log SCC	2.3444	2.1157	2.6911	2.7862	2.6669
sd	0.6839	0.5354	0.7407	0.9419	0.6895
xg	221	131	491	611	464

n = number of cases of individual milk samples; x = arithmetic mean; sd = standard deviation; vx = coefficient of variation in %; xg = geometric mean

III: Test of differences between groups of somatic cell counts (SCC) in milk with different mastitis state with different pathogen occurrence respectively (from Tab. II)

Difference	n	t	significance level
negative versus positive in total	161	5.68	$P < 0.001$
negative versus staphylococci	110	3.76	$P < 0.001$
negative versus other positive	148	5.34	$P < 0.001$
staphylococci versus other positive	64	0.51	$P > 0.05$

t = t-test criterion value



1: Medians of case groups of somatic cell counts (SCC in ths.ml⁻¹) in milk with various mastitis state of mammary gland of animals with different pathogen occurrence respectively

sample groups under presumption of normal frequency distribution of logarithmic transformed SCC values. This limit for suspicion of infection is 159 for positive group, 113 for staphylococci and 174 ths.ml⁻¹ for other positive. This could be possible to recommend the value 174 ths.ml⁻¹ as limit for suspicion of infection and for practical use with target to apply selective mastitis antibiotic treatment (for instance at dairy cow drying) or other preventive or curative measures.

Acknowledgement

This work was supported by projects of Ministry of Agriculture NAZV KUS QJ1210301, QJ1230044, QJ1210284 and NAZV QH81105.

REFERENCES

- ALI, A. K. A., SHOOK, G. E., 1980: An optimum transformation for somatic cells concentration in milk. *J. Dairy Sci.*, 63, 3: 487–490. ISSN 0022-0302.
- ANDREWS, R. J., KITCHEN, B. J., KWEE, W. S., DUNCALFE, F., 1983: Relationship between individual cow somatic cell counts and the mastitis infection status of the udder. *Austral. J. Dairy Techn.*, 38, 2: 71–74. ISSN 0004-9433.
- BENDA, P., VYLEŤELOVÁ, M., TICHÁČEK, A., 1997: A method of prevalence estimation of intramammary *Staphylococcus aureus* and *Streptococcus agalactiae* infection in herds by examination of bulk milk samples. *Vet. Med. – Czech*, 42, 4: 101–109. ISSN 0375-8427.
- BERRY, D. P., O'BRIEN, B., O'CALLAGHAN, E. J., SULLIVAN, K. O., MEANEY, W. J., 2006: Temporal trends in bulk tank somatic cell count and total bacterial count in Irish dairy herds during the past decade. *J. Dairy Sci.*, 89, 10: 4083–4093. ISSN 0022-0302.
- BOŞGELMEZ-TINAZ, G., ULUSOY, S., ARIDOGAN, B., COŞKUN-ARI, F., 2006: Evaluation of different methods to detect oxacillin resistance in *Staphylococcus aureus* and their clinical laboratory utility. *European J. Clin. Microbiol. Inf. Dis.*, 25, 6: 410–412. ISSN 0934-9723.
- BRADLEY, A. J., 2002: Bovine mastitis: An evolving disease. *Vet. J.*, 163, 1: 1–13. ISSN 1090-0233.
- ČSN EN ISO 13366-2 (570531) (2007): *Mléko – Stanovení počtu somatických buněk – Část 2: Návod pro ovládání fluoro-opto-elektronického přístroje*. Praha: ČNI.
- DOHOO, I. R., MEEK, A. H., MARTIN, S. W., BARNUM, D. A. 1981: Use of total and differential somatic cell counts from composite milk samples to detect mastitis in individual cows. *Can. J. Comp. Med.*, 45, 1: 8–14. ISSN 0008-4050.
- EMANUELSON, U., WEVER, P., 1989: Potential of differential somatic cell counts as indicators of mastitis in quarter milk samples from dairy cows. *Acta Vet Scand.*, 30, 4: 475–481. ISSN 0044-605X.
- ERSKINE, R. J., EBERHART, R. J., HUTCHINSON, L. J., SPENCER, S. B., 1987: Herd management and prevalence of mastitis in dairy herds with high and low somatic cell counts. *J. Am. Vet. Med. Assoc.*, 190, 11: 1411–1416. ISSN 0003-1488.
- FRELICH, J., ŠLACHTA, M., 2011: Impact of seasonal grazing on udder health of cows. *Acta. univ. agric. et silvic. Mendel. Brun.*, 59, 1: 53–58. ISSN 1211-8516.
- GOŁĘBIEWSKI, M., BRZOZOWSKI, P., GOŁĘBIEWSKI, L., 2011: Analysis of lactation curves, milk constituents, somatic cell count and urea in milk of cows by the mathematical model of Wood. *Acta Vet. Brno*, 80, 1: 73–80. ISSN 0001-7213.
- HANUŠ, O., TICHÁČEK, A., 1997: Analysis of milking technique effect on somatic cell counts. *Stočarstvo, Animal Husbandry*, 51, 2: 121–128. ISSN 1450-9156.
- HECK, J. M. L., VAN VALENBERG, H. J. F., DIJKSTRA, J., VAN HOOIJDONK, A. C. M., 2009: Seasonal variation in the Dutch bovine raw milk

- composition. *J. Dairy Sci.*, 92, 10: 4745–4755. ISSN 0022–0302.
- HOLMES, M. A., ZADOKS, R. N., 2011: Methicillin resistant *S. aureus* in human and bovine mastitis. *J. Mamm. Gland Biol. Neopl.*, 16, 4: 373–382. ISSN 1535-3702.
- JANŠTOVÁ, B., DRAČKOVÁ, M., DLESKOVÁ, K., CUPÁKOVÁ, Š., NECIDOVÁ, L., NAVRÁTILOVÁ, P., VORLOVÁ, L., 2011: Quality of raw milk from a farm with automatic milking system in Czech Republic. *Acta Vet. Brno*, 80, 2: 207–214. ISSN 0001–7213.
- KALMUS, P., AASMÄE, B., KÄRSSIN, A., ORRO, T., KASK, K., 2011: Udder pathogens and their resistance to antimicrobial agents in dairy cows in Estonia. *Acta Vet. Scand.*, 53, 4: 1–7. ISSN 0044–605X.
- KEEFE, G. P., 1997: Streptococcus agalactiae mastitis: A review. *Can. Vet. J.*, 38, 7: 429–437. ISSN 0008–5286.
- KOVÁČ, G., TÓTHOVÁ, C., NAGY, O., SEIDEL, H., 2011: Milk amyloid A and selected serum proteins in cows suffering from mastitis. *Acta Vet. Brno*, 80, 1: 3–9. ISSN 0001–7213.
- PAŘILOVÁ, M., JEŽKOVÁ, A., STÁDNÍK, L., ŠTOLC, L., 2010: Effect of milking vacuum and overmilking on selected milking characteristics. *Výzkum v chovu skotu / Cattle Research*, LII, 191, 3: 35–43. ISSN 0139–7265.
- PAŘILOVÁ, M., STÁDNÍK, L., JEŽKOVÁ, A., ŠTOLC, L., 2011: Effect of milking vacuum level and overmilking on cows' teat characteristics *Acta univ. agric. et silvic. Mendel. Brun.*, 59, 5: 193–202. ISSN 1211–8516.
- PITKÄLÄ, A., HAVERI, M., PYÖRÄLÄ, S., MYLLYS, V., HONKANEN-BUZALSKI, T., 2004: Bovine mastitis in Finland 2001 – prevalence, distribution of bacteria, and antimicrobial resistance. *J. Dairy Sci.*, 87, 8: 2433–2441. ISSN 0022–0302.
- RAJALA-SCHULTZ, P. J., GRÖHN, Y. T., MCCULLOCH, C. E., GUARD, C. L., 1999: Effects of clinical mastitis on milk yield in dairy cows. *J. Dairy Sci.*, 82, 6: 1213–1220. ISSN 0022–0302.
- RAUBERTAS, J. K., SHOOK, G. E., 1982: Relationship between lactation measures of somatic cell concentration and milk yield. *J. Dairy Sci.*, 65, 3: 419–425. ISSN 0022–0302.
- RENEAU, J. K., 1986: Effective use of dairy herd improvement somatic cell counts in mastitis kontrol. *J. Dairy Sci.*, 69, 6: 1708–1720. ISSN 0022–0302.
- RENEAU, J. K., APPLEMAN, R. D., STEUERNAGEL, G. R., MUDGE, J. W., 1983, 1988: *Somatic cell count. An effective tool in controlling mastitis*. Minnesota: Agricultural Extension Service, University of Minnesota, AG-FO-0447, 1983 and 1988.
- SAWA, A., PIWCZYNSKI, D., 2002: Somatic cell count and milk yield and composition in White × Holstein-Friesian cows. *Med. Weter.*, 58, 8: 636–640. ISSN 0025-8628.
- SHELDRAKE, R. F., HOARE, R. J. T., MCGREGOR, G. D., 1983: Lactation stage, parity, and infection affecting somatic cells, electrical conductivity, and serum albumin in milk. *Austr. J. Dairy Sci.*, 66, 3: 542–547. ISSN 0022–0302.
- TOUŠOVÁ, R., STÁDNÍK, L., DUCHÁČEK, J., BARÁKOVÁ, M., 2011: Influence of vaccination against mastitis on its incidence, somatic cells amount and content of solid components in milk of primiparous dairy cows. *Výzkum v chovu skotu / Cattle Research*, LIII, 196, 4: 84–90. ISSN 0139–7265.
- VASIL M., 2009: Etiology, course and reduction of incidence of environmental mastitis in the herd of dairy cows. *Slovak J. Anim. Sci.*, 42, 3: 136–144. ISSN 1337–9984.
- VYLETĚLOVÁ, M., 2006: Development of mastitis pathogen occurrence since year 1993. In: *Agroregion 2006. VI. ročník mezinárodní konference 24.–25. 8. 2006, Zvyšování konkurenceschopnosti v zemědělství*. České Budějovice: Jihočeská univerzita České Budějovice: 32–35. ISBN 80-7040-869-3.
- VYLETĚLOVÁ, M., HANUŠ, O., KARPÍŠKOVÁ, R., ŠTÁSTKOVÁ, Z., 2011: Occurrence and antimicrobial sensitivity in staphylococci isolated from goat, sheep and cow's milk. *Acta univ. agric. et silvic. Mendel. Brun.*, 59, 3: 209–214. ISSN 1211–8516.
- VYLETĚLOVÁ, M., HANUŠ, O., KUČERA, J., 2009: Relationship of milking equipment type to the expressions of milk secretion disorders. *Acta fytotech. zootech.*, 12, Special Issue: 105–107. ISSN 1335–258X.
- WIGGANS, G. R., SHOOK, G. E., 1987: A lactation measure of somatic cell count. *J. Dairy Sci.*, 70, 12: 2666–2672. ISSN 0022–0302.

Address

RNDr. Marcela Vyleťelová Klimešová, Ph.D., doc. Ing. Oto Hanuš, Ph.D., Ing. Petr Roubal, CSc., Ludmila Nejeschlebová, Dairy Research Institute, Prague, Ke Dvoru 12a, 160 00 Praha 6 - Vokovice, Czech Republic, MVDr. Lucie Hasoňová, Ph.D., University of South Bohemia in České Budějovice, Agricultural Faculty, Studentská 13, 370 05 České Budějovice, Czech Republic, Mgr. Ivan Manga, Ph.D., Agritec Plant Research, Ltd., Zemědělská 16, 787 01 Šumperk, Czech Republic, e-mail: marcela.vyletelova@seznam.cz