

EVALUATION OF DEVELOPMENT IN INDIRECT DETERMINATION OF MILK FAT FREE FATTY ACIDS IN CZECH REPUBLIC

Oto Hanuš, Eva Samková, Jan Říha, Marcela Vyletělová Klimešová, Petr Roubal

Received: May, 2013

Abstract

HANUŠ OTO, SAMKOVÁ EVA, ŘÍHA JAN, VYLETĚLOVÁ KLIMEŠOVÁ MARCELA, ROUBAL PETR: *Evaluation of development in indirect determination of milk fat free fatty acids in Czech Republic*. Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis, 2013, LXI, No. 6, pp. 1669–1679

Free fatty acids (FFAs) in fat are important indicator of raw milk quality. Result reliability of FFAs is important. Aim was to verify MIR–FT (mid infrared spectroscopy with Fourier's transformations) method for its calibration to determine FFAs, time stability of MIR–FT FFA calibration and calibration levelling in laboratory networks. Reference (RE) milk samples (1 set = 8) were prepared according to CSN 57 0533 (FFAs in $\text{mmol} \cdot 100\text{g}^{-1}$ of fat). MIR–FT instruments were: 1 LactoScope FTIR (DE); 2 Bentley FTS (BE); 2 MilkoScan FT 6000 (FO). 3 calibrations of MIR–FT (5) in 3 laboratories were performed. Bulk milk samples came from 4 herds and 2 breeds. These 4 samples were used for calibration in native and modified form. Modification increased FFAs by cca 100%. Calibration set had 8 samples. 1 between calibration interval was checked monthly by proficiency testing (PT). PT set had 10 samples. 5 samples were with native milk and 5 had modified fat content, lower and higher. Maximal value of difference variability for calibration quality validation is x (sd of difference MIR–FT and RE) plus 1.64 multiple of sd (on 95% level), $1.0613 \text{ mmol} \cdot 100\text{g}^{-1}$. Mean validation correlation coefficient (r) between MIR–FT and RE results was 0.802 ± 0.082 ($P < 0.001$), from 0.666 to 0.945. Minimal value at calibration is x minus 1.64 multiple of sd (0.668). Correlations between MIR–FT results were higher by 8.4% (0.869 ($P < 0.001$) > 0.802). Example PT with 10 and 5 milk samples had similar results of r 0.887 and 0.953 ($P < 0.001$ and $P < 0.05$). There is possibility to construct a levelling programme for calibrated instruments. Some equation between PT reference and instrumental values could correct MIR–FT results for their better comparability.

raw cow milk quality, fat, reference, titration, infrared spectroscopy, stability of calibration, result reliability, proficiency testing, result levelling

Determination and interpretation of free fatty acids (FFAs) in milk

Small portion of fatty acids in milk which are not esterified in triglycerides is freely diffused mostly in fat phase and a little bit in water phase and it is called as free fatty acids (FFAs). Current content of FFAs in milk fat lies between 0.5 and 1.2 $\text{mmol} \cdot 100\text{g}^{-1}$ and maximal enabled is 13.0 $\text{mmol} \cdot 100\text{g}^{-1}$ for method by churning and 32.0 $\text{mmol} \cdot \text{kg}^{-1}$ for method by extraction and titration according to CSN 57 0529 and CSN 57 0533 (Cvak *et al.*, 1992). Gerber's acidobutyrometrical method for milk fat determination holds as many as 90% of FFA

content into milk fat portion but on the contrary the extraction gravimetric method according to Röse-Gottlieb does not include FFAs into fat portion so reliably and thus 70% of them is lost in this way (Kerkhoff Mogot *et al.*, 1982).


Increase of FFAs means negative impacts as lipolysis usually from reason of metabolic problems of dairy cow (Fig. 1). Increased concentration of FFAs causes an aggravation of milk technological properties (Vyletělová *et al.*, 2000 a, b) but mostly deterioration of sensory milk properties as taste and flavour. After that it has lightly bitter smack as consequence and this can damage quality of dairy

products. Fat destruction is phenomenon which is caused by native enzymes as lipases (Antonelli *et al.*, 2002; Deeth, 2006; Ferlay *et al.*, 2006) in milk or by lipases which are supplied by bacterial contamination of milk. Therefore lipolysis is spontaneous or induced (Fig. 1). Of course, lipases can be thermoresistent and thus in this way to influence milk also after its heat treatment by dairy product degradation. Wasteful milk handling as often pumping and ripple at manipulation (Sjaunja, 1984; Thomson *et al.*, 2005; Hanuš *et al.*, 2008 b; Genčurová *et al.*, 2009 and 2011) and its freezing on technology surfaces also induce own lipolysis. Heat and mechanical energy which is added into multicomponent milk system destroys the membranes of fat globules and thus releases fatty acids from esteric linkage of triglycerides. Therefore, milk stream (Peterková, 2002) should not exceed the speed 1 till 1.5 m.s⁻¹. Poor hygiene of dairy cow stabling and milking as well as bad storage and treatment of raw milk can lead to propagation of undesirable psychrotrophic, thermoresistant and sporulating milk microflora (Vyleťelová *et al.*, 1999 a, b; Cempírková and Thér, 2000; Cempírková, 2001, 2002, 2007; Dankow *et al.*, 2004; Hanuš *et al.*, 2004; Foltys and Kirchnerová, 2006 and 2010; Hantis-Zacharov and Halpern, 2007; Torkar and Teger, 2008; Cempírková and Mikulová, 2009; Cempírková *et al.*, 2009). The mentioned facts can increase the lipolysis intensity (Fig. 1).

Definition of FFAs and paper goal

FFAs are a mixture of fatty acids released from milk fat by lipolysis or such which overpassed from animal blood and body fat tissues. In terms of proportions this mixture is influenceable by animal nutrition and health state, season and other factors which means that this is very hardly seizable, expressible and interpretable from analytical and molar point of view respectively. Analytically this is result of alkaline titration which is not in constant ratio to molar concentrations of individual fatty acids. However, this way of expression corresponds very good to practical dairy purposes as conventional interpretation. Values of FFAs can serve to control the health of dairy cows or raw milk quality in consideration of quality and shelf-life of resulting milk products.

FFA analytical methods are relatively complicated in terms of reliability and expressin of units in spite of matter definition simplicity. Variable mixture of acids form broken triglycerides is namely instable in composition ratios. Authors of various articles were concerned with FFA analytical methods (Sjaunja, 1984; Koops *et al.*, 1990; Foss, 2001, 2004; Bijgaart, 2006; Hanuš *et al.*, 2008 a, 2009) and above mentioned methodical complications ensued on these papers. Reference and routine FFA analytical methods can be so called extraction-titration method, churning-titration method, BDI and mid infrared spectroscopy (MIR) also in modification with Fourier's transformations (MIR-FT; CSN 57 0533; Koops *et al.*, 1990; IDE, 1991; Cvak *et al.*,



1: Lipolysis rise of fat in raw milk, content increase of free fatty acids (FFA), threat of milk and dairy product quality – factors and their combinations, related to animal and technology (modified according to Sjaunja (1984), Shelley *et al.* (1987), O'Brian *et al.* (1998), Vyleťelová *et al.* (2000 a), Ma *et al.* (2000), Antonelli *et al.* (2002), Santos *et al.* (2003), Wiking *et al.* (2003, 2006), Hanuš *et al.* (2004, 2008 b), Thomson *et al.* (2005), Ferlay *et al.* (2006), Genčurová *et al.* (2009 and 2011) and Mikulová (2011))

1992; Foss, 2001, 2004; Thomson *et al.*, 2005; Bijgaart, 2006; Mikulová, 2011).

Therefore, aim of this paper was to verify possibilities of MIR-FT method in terms of its calibration to milk fat free fatty acids (FFA) determination, time stability of MIR-FT FFA calibration and calibration levelling in instrument working nets of dairy laboratories.

MATERIAL AND METHODS

Analytical and instrumental systems for determination of FFAs

Experimental and development methodical observation was carried out in CR reference and routine milk laboratory network as a case study. One reference laboratory (A) and two routine laboratories (B and C) with one reference method and five MIR-FT instruments (three types) were included in this paper. Calibration (reference) sets of milk samples (one set is equal to eight samples) were prepared according to previously tested procedures (Hanuš *et al.*, 2008 a, b, 2009; Genčurová *et al.*, 2009, 2011) and in accordance with standard CSN 570533 using churning method in reference laboratory. The following MIR-FT instruments with indirect measurement principle were included in this analytical system and its evaluation: once LactoScope FTIR (DE; Delta Instruments, The Netherlands); two times Bentley FTS (BE; Bentley Instruments, USA); two times MilkoScan FT 6000 (FO; Foss, Denmark).

Experimental reference sample preparation, pilot calibration and between calibration intervals

Milk for preparation of calibration samples and for reference method was stored and transported in the same way. Before analyse the milk samples were in water bath only time which was necessary to reach 40 °C for measurement. Whole sample mixing was made carefully. Reference method was carried out two times for calibration samples. Churning-titration method (with KOH titration, CSN 570533) was used as reference (RE) procedure for determination of FFAs and results were expressed in mmol.100g⁻¹ of fat.

During one year three (I, II and III) pilot calibrations (two winter in February and November and one summer in August) of five MIR-FT instruments in three dairy laboratories, which were linked in network, were performed. Bulk milk samples came from four dairy cow herds, two with Czech Fleckvieh and two with Holstein breed. One Holstein herd had poorer energy nutrition and higher native content of FFAs in calibration sample set was established in this way. These four samples were used for calibration in native form and also were stressed by mechanical stirring (Hanuš *et al.*, 2008 a, b, 2009; Genčurová *et al.*, 2009, 2011) for increase of FFAs. One litre of

milk was in vessel (height 19 cm, diameter 13 cm), temperature from 18 to 22 °C, eccentric stirrer (stainless steel surface 7 × 7 cm) with perforation like three circle inlets (diameter 12 mm), 550 wheel revolutions per minute, stirring time from 15 to 20 minutes. It was used in first two calibrations (I and II). This kind of treatment can increase content of FFAs approximately by 100% (60–140%). Reduced milk mechanical stress (10 minutes) was used at last calibration (III). In this way the calibration set consisted regularly of eight milk samples.

One between calibration interval was checked monthly by proficiency testing (PT; Grappin, 1993; Leray, 1993; 2006). PT sample set consisted of ten milk samples for current commercial MIR-FT calibration of main milk components such as fat, protein, casein, lactose and solids non fat content. Five bulk samples were native milk (Czech Fleckvieh and Holstein breed) and five samples had artificially modified fat content to lower (2, milk dilution) and higher (3, cream addition) values.

Calibration (reference) and also PT (control) samples were cooled, preserved (Broad Spectrum Microtabs II, DF Control Systems, England; 0.02%) and sent to laboratories for measurement in thermoboxes under controlled conditions (Sojková *et al.*, 2009). Calibration, analytical work and instrument maintenance were performed according to relevant producer operation manuals.

Statistical procedures

Evaluation of calibration results was performed in accordance with relevant system of statistical treatment (Grappin, 1987, 1993; Hanuš *et al.*, 2008 a, 2009; Genčurová *et al.*, 2011) and proficiency testing was evaluated by Euclidian distance from origin (Leray, 1993; 2006) using Microsoft Excel programme on the basis of difference comparison or testing and regression analyse.

RESULTS AND DISCUSSION

Molecularly inhomogeneous mixture of milk FFAs can be a source of variations in calibration certainty and result reliability of MIR-FT method in dependence on various factors as animal feeding, hygiene and milking considerateness, milk storage and transport. Nevertheless, also the reference method is assessable to these factors in sure sense when in its principle (KOH titration) also grades mentioned FFA molecular variability. MIR-FT use for FFA determination and raw milk quality control is still at the beginning for these various theoretical and practical questions of MIR-FT application, especially in laboratory networks. The development of suitable MIR-FT calibration procedures and developing targeted measures in laboratory networks could contribute to increase of reliability of instrumental FFA determination in milk laboratories.

In Tab. I, II and III there are shown the results of three calibrations of five MIR-FT instruments

(BE1 and 2, FO1 and 2 and DE) for determination of milk FFAs. Means and standard deviations ($\bar{x} \pm \text{sd}$) are mentioned before (FM) and after (CM; as validation result of accepted calibration) calibration. As it is seen according to mutual agreement among reference and instrumental means (CM), the calibration results were the best in first calibration (Tab. I) and the worse in third calibration (Tab. III)

in the laboratory network. Due to lower mechanical stress of chosen milk samples the RE mean of FFAs was lowest ($1.152 \text{ mmol.100g}^{-1}$) in the third calibration (Tab. III) as compared to the first and second (Tab. I and II; 1.934 and $1.746 \text{ mmol.100g}^{-1}$). The real reasons for this fact are unknown. In the same sense, there is also lower variability of FFA values (Tab. III and II; $0.8102 < 1.2215 \text{ mmol.100g}^{-1}$).

I: Results of milk fat FFA calibrations (in mmol.100g^{-1}) of MIR-FT instruments according to reference method results in August 2012

			1	2	3	4	5	6	7	8	x	sd
A	RE		0.870	2.900	0.991	2.335	1.385	2.832	1.823	2.335	1.934	0.7926
A	DE	FM	0.982	1.923	1.005	1.565	1.027	2.113	1.082	2.264	1.495	0.5417
	DE	CM	1.320	2.930	1.440	2.300	1.470	2.630	1.490	3.300	2.110	0.7803
B	BE1	FM	1.585	2.220	1.840	2.235	1.325	2.185	2.125	2.490	2.001	0.3867
	BE1	CM	1.310	2.270	1.690	2.300	0.910	2.220	2.125	2.680	1.938	0.5879
C	BE2	FM	1.898	2.405	1.661	1.856	1.655	2.350	1.960	2.429	2.026	0.3232
	BE2	CM	1.223	2.301	1.392	2.062	1.073	2.425	2.121	2.898	1.937	0.6433
C	FO1	FM	0.511	2.194	0.595	1.630	0.468	2.495	0.805	3.327	1.503	1.0804
	FO1	CM	0.844	2.608	1.034	2.097	0.953	2.966	1.205	3.790	1.937	1.1009
C	FO2	FM	1.214	1.750	1.152	1.468	1.274	1.902	1.285	2.000	1.506	0.3325
	FO2	CM	1.305	2.364	1.326	1.924	1.501	2.699	1.651	3.052	1.978	0.6587


Laboratory A, B and C; RE reference method; DE Delta; BE Bentley; FO Foss; FM first measurement; CM control measurement; 1–8 milk samples; x arithmetical mean; sd standard deviation

II: Results of milk fat FFA calibrations (in mmol.100g^{-1}) of MIR-FT instruments according to reference method results in November 2012

			1	2	3	4	5	6	7	8	x	sd
A	RE		0.782	3.606	1.049	3.398	1.053	2.379	0.430	1.271	1.746	1.2215
A	DE	FM	1.409	3.295	1.395	3.069	1.556	4.273	1.147	2.063	2.276	1.1352
	DE	CM	0.970	2.480	1.290	2.400	1.280	3.530	0.890	1.620	1.808	0.9177
B	BE1	FM	0.710	2.170	0.130	1.500	0.550	1.830	0.000	1.025	0.989	0.7884
	BE1	CM	1.200	2.890	1.750	2.190	1.250	3.300	0.540	1.480	1.825	0.9212
C	BE2	FM	1.425	2.937	1.314	2.627	1.260	2.373	0.234	2.728	1.862	0.9454
	BE2	CM	1.429	3.343	0.862	3.099	1.353	2.416	0.444	2.080	1.878	1.0375
C	FO1	FM	1.161	3.426	1.262	3.591	1.386	5.675	0.794	2.103	2.425	1.6783
	FO1	CM	0.973	3.132	1.131	3.258	1.248	5.458	0.593	1.809	2.200	1.6428
C	FO2	FM	1.540	2.905	1.282	3.291	2.083	5.809	0.052	1.274	2.280	1.7494
	FO2	CM	1.540	2.869	1.282	3.257	2.078	5.766	0.052	1.203	2.256	1.7385

III: Results of milk fat FFA calibrations (in mmol.100g^{-1}) of MIR-FT instruments according to reference method results in February 2013

			1	2	3	4	5	6	7	8	x	sd
A	RE		0.649	2.396	0.701	2.049	0.180	0.660	0.761	1.821	1.152	0.8102
A	DE	FM	0.460	1.344	0.617	1.336	0.496	0.978	0.661	2.166	1.007	0.5853
	DE	CM	0.710	1.530	0.830	1.620	0.670	1.170	0.880	2.180	1.199	0.5354
B	BE1	FM	0.000	1.135	0.055	0.950	0.000	0.510	0.495	1.255	0.550	0.5146
	BE1	CM	0.000	0.860	0.600	1.540	0.000	1.090	0.670	1.580	0.793	0.6071
C	BE2	FM	0.657	0.786	0.681	0.762	0.651	0.805	0.750	0.742	0.729	0.0589
	BE2	CM	0.666	1.823	0.611	1.607	0.060	1.091	0.885	1.718	1.057	0.6215
C	FO1	FM	0.995	2.078	1.039	2.057	0.567	1.877	2.975	4.072	1.958	1.1489
	FO1	CM	0.409	0.848	0.423	0.837	0.211	0.704	1.184	1.584	0.775	0.4491
C	FO2	FM	0.661	1.882	0.906	1.736	0.655	1.331	1.263	2.800	1.404	0.7244
	FO2	CM	0.271	0.764	0.367	0.703	0.245	0.495	0.500	1.064	0.551	0.2786



2: Closest and weakest regression relationship of determination of FFAs (in mmol.100g⁻¹) between RE (x) and MIR-FT (y) as calibration validation ($r=0.945$ and 0.666 , $P < 0.001$ and $P > 0.05$)


From this results (Tab. I, II and III), there is possible to derive an maximal value as statistical parameter of calibration quality for its validation under practical laboratory conditions. This is x (sd of d, mean standard deviation of difference means between MIR-FT and RE) plus 1.64 (one side limitation) multiple of relevant sd (on 95% level of certainty, according to Grappin, 1987), which is $1.0613 \text{ mmol.100g}^{-1}$ ($0.622474 + 0.2662 \times 1.64$). This is visibly higher value than estimated Genčurová *et al.* (2011) at smaller data file, $0.75 \text{ mmol.100g}^{-1}$.

IV: Validation correlation coefficients between instrumental (MIR-FT) and RE results of FFAs in three calibrations (CM, after calibration)

Calibration	I	II	III
DE x RE	0.856 ***	0.784 *	0.827 *
BE1 x RE	0.755 *	0.816 *	0.708 ns
BE2 x RE	0.827 *	0.945 ***	0.938 ***
FO1 x RE	0.821 *	0.743 *	0.683 ns
FO2 x RE	0.837 ***	0.666 ns	0.831 ***
x	0.819	0.791	0.797
sd	0.038	0.103	0.104

Statistical significance: *** $P \leq 0.001$; ** $P \leq 0.01$; * $P \leq 0.05$; ns (no significance) $P > 0.05$

The best relationships (correlation coefficients (r) 0.856, 0.945 and 0.837; $P < 0.001$) from three calibrations between MIR-FT instrumental determination (according to instruments DE, BE and FO) of FFAs and results of reference method (RE) after calibration (CM; validation correlation coefficients) are shown in Tab. IV and Fig. 2 as example. According to instruments their statistical parameters as $x \pm \text{sd}$, minimum and maximum were (Tab. IV): for DE 0.822 ± 0.036 , from 0.784 to 0.856; for BE 0.832 ± 0.095 , from 0.708 to 0.945; for FO 0.764 ± 0.077 , from 0.666 to 0.837. In terms of total statistic these were: 0.802 ± 0.082 ($P < 0.001$), from 0.666 to 0.945. It means that on the average 64.3% (89.3% as maximum) of variability in MIR-FT FFA results could be explained by variability in the real FFA results using reference method. From this results, there is possible to derive an minimal value as statistical parameter of calibration quality for its validation under practical laboratory conditions. This is x minus 1.64 multiple of sd (on 95% level of certainty, according to Grappin (1987), one side limitation), which is 0.668 (minimal acceptable r). This is lower value than estimated Genčurová *et al.* (2011) at smaller data file, 0.841. Foss professional materials (2001, 2004) show similar possibilities with r 0.897 as one regular example ($n = 1.927$). General linear regression equation for all results of



3: General regression of relationship of FFA results (in mmol.100g⁻¹) between MIR-FT (x, 5 instruments) and reference method (y) at calibration I ($r=0.784$; $P < 0.01$)

V: Correlation coefficients between instrumental (MIR-FT) results of FFAs in three calibrations (CM, after calibration)



Calibration	I	II	III
DE-BE1	0.802 *	0.943 ***	0.887 ***
DE-FO1	0.977 ***	0.995 ***	0.786 *
DE-FO2	0.955 ***	0.937 ***	0.979 ***
DE-BE2	0.879 ***	0.764 *	0.889 ***
BE1-FO1	0.824 *	0.923 ***	0.745 *
BE1-FO2	0.783 *	0.893 ***	0.863 ***
BE2-FO1	0.915 ***	0.735 *	0.724 *
BE2-FO2	0.912 ***	0.662 ns	0.899 ***
BE1-BE2	0.953 ***	0.782 *	0.833 ***
FO1-FO2	0.988 ***	0.956 ***	0.871 ***
x	0.899	0.859	0.848
sd	0.074	0.113	0.077

instruments (x axis) in the network in relation to RE values (y axis) in one calibration (I) is shown in Fig. 3 as example ($r = 0.784$; $P < 0.01$). Of course, this kind of evaluation could be probably useable as model for levelling of results of all instruments in good technical condition in laboratory network during following between calibration interval.



In Tab. V, there are shown the correlations (r) between instrumental calibration FFA results mutually. Only one result is insignificant ($P > 0.05$) and highest coefficient is 0.995 ($P < 0.001$; Fig. 4). In general, these coefficients are higher as compared to those in Tab. IV (MIR-FT to RE values) by 8.4% on

the average (mean r 0.869 ($P < 0.001$) > 0.802) which is probably caused by higher methodical homogeneity (Tab. V) in this file. Further, interesting thing is that these coefficients are usually higher between the same types of instruments (in total 0.894 ± 0.079 , from 0.782 to 0.988 (BE x BE and FO x FO), $n = 6$, where BE 0.856 ± 0.088 and FO 0.938 ± 0.06) than between different instruments (0.861 ± 0.029 , from 0.662 to 0.995 (DE x BE, DE x FO and BE x FO), $n = 24$). This fact shows logically on various solutions of infrared signal processing and its information recovery between used instrument types. However, from analytical point of view, this is possible also correlations between different kinds of instruments to specify as suitable. Also, as it has been mentioned already these „hetero- and homo-methodical“ (Tab. IV (0.802, RE and MIR-FT) and V (0.869, MIR-FT)) correlation coefficients between FFA results (as minor milk component) are still lower than the same type of correlations at investigation of major milk components such as fat, protein, lactose and solid non fat (RE x MIR-FT and inside MIR-FT) in accordance with other professional materials (Koops *et al.*, 1990; Foss, 2004; Bijgaart, 2006; Hanuš *et al.*, 2008 a, 2009; Genčurová *et al.*, 2011).

Example results from one PT about determination of FFAs are shown in Fig. 5 and 6 as example. These were carried out for ten milk calibration samples during current milk composition MIR-FT calibration (fat, protein, lactose, solid non fat) including fat content modified samples (Fig. 6). It means that no all samples have native composition




4: Closest and weakest regression relationship of determination of FFAs (in mmol.100g⁻¹) between MIR-FT mutually after carried out calibration ($r = 0.995$ and 0.662 , $P < 0.001$ and $P > 0.05$)



5: Example of regression relationship of determination of FFAs (in mmol.100g⁻¹) between MIR-FT mutually in proficiency testing for 10 (native and modified bulk) and 5 (native bulk) milk samples ($r = 0.887$ and 0.953 , $P < 0.001$ and $P < 0.05$)

and this test is not regular for PT of FFAs in this way. However, for the future, it could be probably more suitable to perform this PT evaluation only with native bulk milk samples (5 in this case and used system (Fig. 6)) because of result protection against possible interference impacts due to too high variability of main milk components. That is also reason why the PT result pattern can be a little bit different for the same PT trial (Fig. 6). The relationships in Fig. 5 are similar to those in Tab. V for 10 (native and modified) and also 5 (only native) samples by their character. The instrumental r were

quite comparable for the same PT case (Fig. 6), 0.887 and 0.953 ($P < 0.001$ and $P < 0.05$; Fig. 5). Therefore that fact is very interesting that differences among instruments (MIR-FT) in inter-calibration interval are very often stable in terms of trend. It offers us possibility to construct a levelling programme for already calibrated instruments because also mutual correlations are usually close. So some generally calculated regression equation between derived PT reference and instrumental values could correct following MIR-FT results in the network for their better comparability. According to above mentioned



6: Example of evaluation of Euclidian distance from origin for determination of FFAs in PT between MIR-FT for 10 and 5 milk samples

d = mean difference (indirect method – reference value); sd = standard deviation of individual differences along samples; lines which go out from central position of graph represent statistical significance on level 5% (pair t -test, Student's distribution), points which are below lines are significantly different ($P \leq 0.05$), points over lines are insignificantly different ($P > 0.05$); bow incloses 90% of confidence interval of Euclidian distance and successful result in PT, box incloses very successful result.

facts it could be probably carried out mostly in bias field of stated instrumental equations.

In other factors of mentioned problem Thomson *et al.* (2005) found an impact of timing (at sampling and 24 hours after sampling at cold storage) of milk sample preservation (0.1 ml of 0.2% hydrogen peroxide added to 30 ml of milk) on FFA value estimation using BDI method. In this context so called BDI method, which is often used for instance in the Netherlands, serves systematically lightly lower results as compared to reference churning method (CSN 570533) because certain portion of FFAs penetrates into water phase during this procedure (Cvak *et al.*, 1992), so there are principle reasons for such phenomenon. Concentration of FFAs (Thomson *et al.*, 2005) was increased by 21% at lengthening of interval from collection to analyse by 24 hours and by 9% at delayed milk collection from daily to every second day. However the between-farm variation remained reasonably consistent.

Because of preparation way (mechanical stress in some cases) the FFA reference milk samples (Hanuš *et al.*, 2008 a, b, 2009; Genčurová *et al.*, 2009, 2011) are very sensitive to way of treatment and transport. In consideration of these our results, this could be advantageous to reduce used time (approximately by 50%) of mechanic stress at preparation of relevant reference samples because of improvement of stability and shelf-life of these milk FFA calibration samples. That is also reason why it is necessary to use quick and cold it means controlled sample set transport in case of sending a delivery to routine laboratories to calibration in the working

net, for instance under conditions which were experimentally validated by Sojková *et al.* (2009).

CONCLUSION

From analyse of calibration stability variations of MIR-FT method for FFA measurement it is possible to deduce as suitable to construct PC programme for calculation of levelling equations of individual instruments in laboratory working net for improvement of FFA result reliability: – current calibration of instruments on milk FFA measurement will be carried out in the working net; – first possibility is direct use of reference method values from reference sample set after performed calibration for next calculation (a); – second variant is that Grubbs test of remoteness (on level of probability 0.05%) will be used at measurement of reference sample set during actual calibration for insertion of instrumental sample results into calculation of reference sample value (b); – this value will be created by arithmetical mean of purged set of sample values according to individual instruments (laboratories); – sample values of reference set of each instrument will be related to relevant reference values of instrument set in working net (a or b); – after that the individual levelling equation will be calculated for each instrument; – this equation will be used during period from actual calibration to next calibration for FFA result transformation at relevant instrument on levelling results of working net.

SUMMARY

Content of free fatty acids (FFAs) in fat is important indicator of raw milk quality in terms of sensorial properties and its product shelf-life. FFAs are product of dairy cow energy metabolism and milk fat lipolysis induced by native or bacterial enzymes. Therefore, their content is influenceable by various factors as animal feeding, hygiene and milking considerateness, milk storage and transport. Such are the reasons why result reliability of analytical methods for series determination of FFAs is important. Therefore, aim of this paper was to verify possibilities of MIR-FT (mid infrared spectroscopy with Fourier's transformations) method in terms of its calibration to determination of milk fat FFAs, time stability of MIR-FT FFA calibration and calibration levelling in instrument networks of dairy laboratories. Reference (RE) milk samples (one set is equal to eight samples) were prepared according to previous procedures and CSN 570533 (FFAs in $\text{mmol} \cdot 100\text{g}^{-1}$ of fat). The following MIR-FT instruments were included: 1 LactoScope FTIR (DE); 2 Bentley FTS (BE); 2 MilkoScan FT 6000 (FO). Three pilot calibrations (two winter in February and November and one summer in August) of five MIR-FT instruments in three dairy laboratories in network were performed. Bulk milk samples came from four dairy cow herds and two breeds. These four samples were used for calibration in native and modified form in terms of fat content. Modification increased FFAs approximately by 100%. Calibration set consisted of 8 milk samples. One between calibration interval was checked monthly by proficiency testing (PT). PT sample set consisted of 10 milk samples. Five bulk samples were native milk and five had artificially modified fat content to lower and higher values. Calibration, analytical work and instrument maintenance were performed according to relevant producer operation manuals. Statistical evaluation procedures of calibration results and PT were on the basis of regression analyse and difference comparison or testing. The calibration results were the best in first calibration and the worse in third calibration in the laboratory network. Due to lower mechanical stress of chosen milk samples the RE mean of FFAs was lowest ($1.152 \text{ mmol} \cdot 100\text{g}^{-1}$) in the third calibration as compared to the first and second (1.934 and $1.746 \text{ mmol} \cdot 100\text{g}^{-1}$). Maximal value of difference variability as validation parameter of calibration quality is x (sd of difference MIR-FT and RE) plus 1.64 multiple of sd (on 95%

level) which is $1.0613 \text{ mmol} \cdot 100\text{g}^{-1}$. Mean validation correlation coefficient (r) between MIR–FT and RE results was 0.802 ± 0.082 ($P < 0.001$), from 0.666 to 0.945. On the average 64.3% (89.3% as maximum) of variability in MIR–FT FFA results could be explained by variability in the RE results. Minimal value as validation parameter of calibration quality is x minus 1.64 multiple of sd which is 0.668. Correlations between MIR–FT results were higher by 8.4% (0.869 ($P < 0.001$) > 0.802) as compared to validation r . Example PT with 10 (native and modified) and 5 (native bulk samples) milk samples had similar results of r 0.887 and 0.953 ($P < 0.001$ and $P < 0.05$). Differences among instruments (MIR–FT) in inter–calibration interval are very often stable. It offers the possibility to construct a levelling programme for calibrated instruments. So some generally calculated regression equation between derived PT reference and instrumental values could correct following MIR–FT results for their better comparability in the relevant network. According to above mentioned facts it could be probably carried out mostly in bias field of stated instrumental equations. According to the obtained results a philosophy of levelling programme was defined in steps as paper conclusion.

Acknowledgement

This paper was supported by projects Ministry of Agriculture No NAZV KUS QJ1210301, RO0513 from February 2013 and NAZV QH81105 and Ministry of Education, Youth and Sports No MSM 6007665806. Authors also thank to Mr. Jaroslav Kopecký and Mrs. Radoslava Jedelská and Mr. Pavel Kopuncz, Zdeněk Motýčka, Jan Zlatníček, Miloš Klimeš and Mrs. Marta Kleinová from Czech–Moravia Breeders Corporation for their kind support and technical cooperation.

REFERENCES

- ANTONELLI, M. L., CURINI, R., SCRICCILO, D., VINCI, G., 2002: Determination of free fatty acids and lipase activity in milk: quality and storage markers. *Talanta*, 58: 561–568. ISSN 0039–9140.
- BIJGAART VAN DEN, H., 2006: New applications of mid-infra-red spectrometry for the analysis of milk and milk products. 2 Free fatty acids. *IDF Bulletin*, 406: 22–28. ISSN 0250–5118.
- CEMPÍRKOVÁ, R., 2007: Contamination of cow's raw milk by psychrotrophic and mesophilic microflora in relation to selected factor. *Czech J. Anim. Sci.*, 52, 11: 387–393. ISSN 1212–1819.
- CEMPÍRKOVÁ, R., 2002: Psychrotrophic vs. total bacterial counts in bulk milk samples. *Vet. Med.–Czech*, 47, 8: 227–233. ISSN 0375–8427.
- CEMPÍRKOVÁ, R., 2001: The technology of dairy cows breeding in relation to the hygienic quality of milk. (In Czech) *Coll. Sci. Pap. Fac. Agric. in České Budějovice, Ser. Anim. Sci.*, 18, 1: 33–46. ISSN 1212–558X.
- CEMPÍRKOVÁ, R., MIKULOVÁ, M., 2009: Incidence of psychrotrophic lipolytic bacteria in cow's raw milk. *Czech J. Anim. Sci.*, 54, 2: 65–73. ISSN 1212–1819.
- CEMPÍRKOVÁ, R., MIKULOVÁ, M., TRÁVNÍČEK, J., 2009: Counts of psychrotrophic lipolytic bacteria in cow's raw milk samples from the aspect of technological quality. *J. Agrobiol.*, 26, 2: 113–121. ISSN 1803–4403.
- CEMPÍRKOVÁ, R., THÉR, R., 2000: The effect of living conditions on selected indexes of raw cow's milk. (In Czech) *Coll. Sci. Pap. Fac. Agric. in České Budějovice, Ser. Anim. Sci.*, 17, 1: 55–71. ISSN 1212–558X.
- CVAK, Z., PETERKOVÁ, L., ČERNÁ, E., 1992: Chemical and physic–chemical methods in milk and dairy product quality control. (In Czech). Praha: VÚP, Milcom servis a.s. Praha, (118) 222. ISBN 80–85120–36–4.
- CSN 57 0529, 1993: Raw cow milk for dairy factory treatment and processing. Praha, ČNI.
- CSN 57 0533, 1997: Milk – determination of matter content of free fatty acids. ČNI Prague.
- DANKOW, R., WOJTOWSKI, J., FAHR, R. D., 2004: Hygienic quality of raw milk in relation to methods of production and storage. *Med. Wet.*, 60: 46–49. ISSN 0025–8628.
- DEETH, H. C., 2006: Lipoprotein lipase and lipolysis in milk. *Inter. Dairy J.*, 16: 555–562. ISSN 0958–6946.
- FERLAY, A., MARTIN, B., PRADEL, P., COULON, J. B., CHILLIARD, Y., 2006: Influence of grass–based diets on milk fatty acid composition and milk lipolytic system in Tarentaise and Montbeliarde cow breeds. *J. Dairy Sci.*, 89: 4026–4041. ISSN 0022–0302.
- FOLTYS, V., KIRCHNEROVÁ, K., 2006: Mesophilic and psychrotrophic aerobe sporulating microorganisms in raw cow's milk. *Centr. Eur. J. Biol.*, 1: 545–560. ISSN 1895–104X.
- FOLTYS, V., KIRCHNEROVÁ, K., 2010: Occurrence and detection of aerobic sporulating microorganisms in raw cow's milk. *Slovak J. Anim. Sci.*, 43, 2: 83–87. ISSN 1337–9984.
- FOSS, 2001: MilkoScan FT 6000; Free fatty acids measurement (FFA). Spectrum Calibration.
- FOSS, 2004: MilkoScan FT 120; Improved milk calibration. Analytical Application Note No. 128e.
- GENČUROVÁ, V., HANUŠ, O., JEDELSKÁ, R., KOPECKÝ, J., 2011: Creation and development of the calibration model for the indirect determination of milk fat free fatty acids and the method of a reference sample preparation. (In Czech) *Výzkum v chovu skotu / Cattle Research*, LIII, 193, 1: 10–20. ISSN 0139–7265.

- GENČUROVÁ, V., HANUŠ, O., KUČERA, J., SOJKOVÁ, K., KOPECKÝ, J., JEDELSKÁ, R., 2009: Free fatty acids as possible indicator for milk quality improvement. *Acta fytotech. zootech.*, 12: 25–28. ISSN 1335–258X.
- GRAPPIN, R., 1987: Definition and evaluation of the overall accuracy of indirect methods of milk analysis – application to calibration procedure and quality control in dairy laboratory. *Bulletin of the International Dairy Federation*, Doc. 208, IDF Provisional Standard 128: 3–12. ISSN 0250–5118.
- GRAPPIN, R., 1993: European network of dairy laboratories. In: *Proceedings of an International Analytical Quality Assurance and Good Laboratory Practice in Dairy Laboratories*. Sonthofen / Germany: 205–211, 1992–05–18/20, Brussels.
- HANTIS–ZACHAROV, E., HALPERN, M., 2007: Culturable psychrotrophic bacteria communities in raw milk and their proteolytic and lipolytic traits. *Appl. Environ. Microbiol.*, 73: 7162–7168. ISSN 0099–2240.
- HANUŠ, O., FRELICH, J., VYLETĚLOVÁ, M., ROUBAL, P., VORLÍČEK, Z., JEDELSKÁ, R., 2004: Technologically difficult, pathogenic and food risky bacterial contamination of raw milk and other materials from dairy cow herds. *Czech J. Anim. Sci.*, 49, 11: 489–499. ISSN 1212–1819.
- HANUŠ, O., GENČUROVÁ, V., KOPECKÝ, J., JEDELSKÁ, R., MOTYČKA, Z., ČERNOCKÝ, M., 2008 a: The interpretation of experimental calibration results of routine IR instruments for determination of free fatty acid contents of milk fat. (In Czech) *Výzkum v chovu skotu / Cattle Research*, L, 182, 2: 55–62. ISSN 0139–7265.
- HANUŠ, O., KUČERA, J., GENČUROVÁ, V., KOPECKÝ, J., JEDELSKÁ, R., 2009: Selected parameters of MIR-FT method validation to measurement of free fatty acids in milk fat for milk quality laboratories. (In Czech) *Výzkum v chovu skotu / Cattle Research*, LI, 187, 3: 27–34. ISSN 0139–7265.
- HANUŠ, O., VEGRICHT, J., FRELICH, J., MACEK, A., BJELKA, M., LOUDA, F., JANŮ, L., 2008 b: Analyse of raw cow milk quality according to free fatty acids contents in the Czech Republic. *Czech J. Anim. Sci.*, 53, 1: 17–30. ISSN 1212–1819.
- IDF, 1991: Determination of free fatty acids in milk and milk products. IDF Bulletin No. 256, Brussels, International Dairy Federation.
- KERKHOF MOGOT, M. F., KOOPS, J., NEETER, R., SLANGEN, K. J., HEMERT VAN, H., KOOYMAN, O., WOOLDRIK, H., 1982: Routine testing of farm tank milk with the Milko–Scan 203. 1. Calibration procedure and small-scale experiments. *Neth. Milk Dairy J.*, 36: 115–133. ISSN 0958–6946.
- KOOPS, J., KLOMP, H., HEMERT VAN, H., 1990: Rapid enzymatic assay of free fatty acids (lipolysis) in farm tank milk by a segmented continuous-flow method. Comparison of the results with those obtained by the BDI procedure. *Neth. Milk Dairy J.*, 44: 3–19. ISSN 0958–6946.
- LERAY, O., 1993: CECALAIT: an organization to support analytical quality assurance in dairy laboratories. *Proceedings of an International Analytical Quality Assurance and Good Laboratory Practice in Dairy Laboratories*. Sonthofen / Germany: 349–360, 1992–05–18/20, Brussels.
- LERAY, O., 2006: Reference and calibration system for routine milk testing – advantages / disadvantages, choice criteria. 3rd ICAR reference laboratory network meeting – Kuopio, Finland – 6th June, 49–65. Breeding, production recording, health and the evaluation of farm animals. EAAP publication No. 121, 2007, *Proceedings of the 35th biennial session of ICAR*: 311–317. ISBN 978–90–8686–030–2.
- MA, Y., RYAN, C., BARBANO, D. M., GALTON, D. M., RUDAN, M. A., BOOR, K. J., 2000: Effects of somatic cell count on quality and shelf-life of pasteurized fluid milk. *J. Dairy Sci.*, 83: 264–274. ISSN 0022–0302.
- MIKULOVÁ, M., 2011: Content of free fatty acids lipolytic bacteria and somatic cells in relation to milking technology. *J. Agrobiol.*, 28, 1: 49–54. ISSN 1803–4403.
- O'BRIAN, B., O'CALLAGHAN, E., DILLON, P., 1998: Effect of milking machine systems and components on free fatty acid levels in milk. *J. Dairy Res.*, 65: 335–339. ISSN 1469–7629.
- PETERKOVÁ, L., 2002: Free fatty acids ratio in milk, factors which affect their concentration and possibilities their determination. (In Czech) *Problematika prvovýroby mléka XXVI*, Medlov, Milcom a. s., Praha, 26–30.
- SANTOS, M. V., MA, Y., BARBANO, D. M., 2003: Effect of somatic cell count on proteolysis and lipolysis in pasteurized fluid milk during shelf-life storage. *J. Dairy Sci.*, 86: 2491–2503. ISSN 0022–0302.
- THOMSON, N. A., WOOLFORD, W. M., COPEMAN, A. P. J., 2005: Milk harvesting and cow factors influencing seasonal variation in the levels of free fatty acids in milk from Waikato dairy herds. *New Zeal. J. Agr. Res.*, 48: 11–21. ISSN 0028–8233.
- TORKAR, K. G., TEGER, S. G., 2008: The microbiological quality of raw milk after introducing the two day's milk collecting system. *Acta Agric. Slov.*, 92: 61–74. ISSN 1854–4800.
- SHELLEY, A. W., DEETH, H. C., MACRAE, I. C., 1987: A numerical taxonomic study of psychrotrophic bacteria associated with lipolytic spoilage of raw milk. *J. Appl. Bacteriol.*, 62: 197–207. ISSN 0021–8847.
- SJAUNJA, L. O., 1984: Studies on milk analysis of individual cow milk samples. III. The effect of different treatments on infrared analyses. *Acta Agric. Scand.*, 34: 273–285. ISSN 0906–4710.
- SOJKOVÁ, K., HANUŠ, O., KOPECKÝ, J., JEDELSKÁ, R., 2009: Determination of thermogradient for interlaboratory milk sample transport. (In Czech) *Výzkum v chovu skotu / Cattle Research*, LI, 187, 3: 35–41. ISSN 0139–7265.
- VYLETĚLOVÁ, M., BENDA, P., HANUŠ, O., KOPUNECZ, P., 1999 a: Determination of total psychrotrophic microorganisms in bulk milk

- samples and their relationship to total count of mikroorganisms. (In Czech) *Czech J. Food Sci.*, 17, 6: 216–222. ISSN 1212–1800.
- VYLETĚLOVÁ, M., FICNAR, J., HANUŠ, O., 2000 a: Effects of lipolytic enzymes *Pseudomonas fluorescens* on liberation of fatty acids from milk fat. *Czech J. Food Sci.*, 18, 5: 175–182. ISSN 1212–1800.
- VYLETĚLOVÁ, M., HANUŠ, O., URBANOVÁ, E., 1999 b: The occurrence of proteolytical and lipolytical bacteria in cow bulk milk samples. (In Czech) *Veterinářství*, 49, 11: 480–482. ISSN 0506–8231.
- VYLETĚLOVÁ, M., HANUŠ, O., URBANOVÁ, E., KOPUNECZ, P., 2000 b: The occurrence and identification of psychrotrophic bacteria with proteolytic and lipolytic activity in bulk milk samples at storage in primary production conditions. (In Czech) *Czech J. Anim. Sci.*, 45: 373–383. ISSN 1212–1819.
- WIKING, L., NIELSEN, J. H., BAVIUS, A. K., EDVARDSSON, A., SVENNERSTEN-SJAUNJA, K., 2006: Impact of milking frequencies on the level of free fatty acids in milk, fat globule size, and fatty acid composition. *J. Dairy Sci.*, 89: 1004–1009. ISSN 0022–0302.
- WIKING, L., BJÖRCK, L., NIELSEN, J. H., 2003: Influence of feed composition on stability of fat globules during pumping of raw milk. *Inter. Dairy J.*, 13: 797–803. ISSN 0958–6946.

Address

prof. Ing. Oto Hanuš, Ph.D., Mgr. Jan Říha, Ph.D., Ing. Petr Roubal, CSc., RNDr. Marcela Vyleťelová Klimešová, Ph.D., Dairy Research Institute, Ltd., Prague, Ke Dvoru 12a, 160 00 Praha 6 - Vokovice, Czech Republic; doc. Ing. Eva Samková, Ph.D., University of South Bohemia České Budějovice, Branišovská 1, 370 05 České Budějovice, Czech Republic, e-mail: hanus.oto@seznam.cz